

OPSONIC AND PROTECTIVE
MONOCLONAL AND CHIMERIC ANTIBODIES
SPECIFIC FOR LIPOTEICHOIC ACID OF GRAM POSITIVE BACTERIA

Inventors:

GERALD W. FISCHER

RICHARD F. SCHUMAN

HING WONG

and

JEFFREY R. STINSON

099915-00901
TOP SECRET 4419660

[illegible][illegible][illegible][illegible][illegible][illegible]

Nonetheless, the battle has not been won. Undoubtedly a significant part of the problem is that bacteria are the product of nearly 3 billion years of natural selection from which they have emerged as an immensely diverse group of organisms that colonize almost all parts of the world and its inhabitants. To begin to understand bacteria requires categorization, and the most fundamental categories for bacteria are their response to the Gram stain, yielding (for the most part) Gram positive bacteria and Gram negative bacteria.

The difference in response to the Gram stain results from differences in bacterial cell walls. The cells walls of Gram negative bacteria are made up of a unique outer membrane of two opposing phospholipid-protein leaflets, with an ordinary phospholipid in the inner leaflet but the extremely toxic lipopolysaccharide in the outer leaflet. The cell walls of Gram positive bacteria seem much simpler in comparison, containing two major components, peptidoglycan and teichoic acids plus additional carbohydrates and proteins depending on the species.

Of the Gram positive bacteria, one of the most common genera is *Staphylococcus*. Staphylococci commonly colonize humans and animals and are an important cause of human morbidity and mortality, particularly in hospitalized patients. Staphylococci are prevalent on the skin and mucosal linings and, accordingly, are ideally situated to produce both localized and systemic infections.

There are two main groups of Staphylococci divided according to the production of "coagulase," an enzyme that causes fibrin to coagulate and to form a clot: coagulase positive and coagulase negative. The coagulase positive *Staphylococcus* species most

frequently pathogenic in humans is *Staphylococcus aureus*. *S. aureus* is the most virulent *Staphylococcus* and produces severe and often fatal disease in both normal and immunocompromised hosts. *Staphylococcus epidermidis* is the most common coagulase negative species.

In recent years, *S. epidermidis* has become a major cause of nosocomial infection in patients whose treatments include the placement of foreign objects such as cerebrospinal fluid shunts, cardiac valves, vascular catheters, joint prostheses, and other implants into the body. *S. epidermidis* and *S. aureus* are common causes of post-operative wound infections and *S. epidermidis* has also become a common cause of peritonitis in patients with continuous ambulatory peritoneal dialysis. In a similar manner, patients with impaired immunity and those receiving parenteral nutrition through central venous catheters are at high risk for developing *S. epidermidis* sepsis. (C.C. Patrick, J. Pediatr., 116:497 (1990)). *S. epidermidis* is now recognized as a common cause of neonatal nosocomial sepsis. Infections frequently occur in premature infants that have received parenteral nutrition which can be a direct or indirect source of contamination.

Staphylococcal infections are difficult to treat for a variety of reasons. Resistance to antibiotics is common and becoming more so. See L. Garrett, The Coming Plague, "The Revenge of the Germs or Just Keep Inventing New Drugs" Ch. 13, pgs. 411-456, Farrar, Straus and Giroux, NY, Eds. (1994). In one study, the majority of Staphylococci isolated from blood cultures of septic infants were multiply resistant to antibiotics (A. Fleer et al., Pediatr. Infect. Dis. 2:426 (1983)). A more recent

study describes methicillin-resistant *S. aureus* (J. Romero-Vivas, et al., Clin. Infect. Dis. 21:1417-23 (1995)) and a recent review notes that the emergence of antibiotic resistance among clinical isolates makes treatment difficult (J. Lee., Trends in Micro. 4(4):162-66 (April 1996). Recent reports in the popular press also describe troubling incidents of antibiotic resistance. See The Washington Post "Microbe in Hospital Infections Show Resistance to Antibiotics," May 29, 1997; The Washington Times, "Deadly bacteria outwits antibiotics," May 29, 1997.

In addition, host resistance to Staphylococcal infections is not clearly understood. Opsonic antibodies have been proposed to prevent or treat Staphylococcal infections. See U.S. Patent No. 5,571,511 to G.W. Fischer issued November 5, 1996, specifically incorporated by reference. The microbial targets for these antibodies have been capsular polysaccharides or surface proteins. As to capsular polysaccharides, the immunization studies of Fattom et al., J. Clin. Micro. 30(12):3270-3273 (1992) demonstrated that opsonization was related to *S. epidermidis* type-specific anti-capsular antibody, suggesting that *S. epidermidis* and *S. aureas* have a similar pathogenesis and opsonic requirement as other encapsulated Gram positive cocci such as *Streptococcus pneumonia*. As to surface proteins, Timmerman, et al., J. Med. Micro. 35:65-71 (1991) identified a surface protein of *S. epidermidis* that was opsonic for the homologous strain used for immunization and for monoclonal antibody production. While other monoclonal antibodies were identified that bound to non-homologous *S. epidermidis* strains, only the monoclonal antibody produced to the homologous strain was opsonic and opsonization was enhanced only to the

homologous strain but not to heterologous strains. Accordingly, based on the studies of Fattom et al., and Timmerman et al., and others in the field (and in contrast to our own studies), one would not expect that an antibody that is broadly reactive to multiple strains of *S. epidermidis* and to *S. aureus* would have opsonic activity against both. This is particularly true for antibodies that bind to both coagulase positive and coagulase negative Staphylococci.

Accordingly, there is a need in the art to provide monoclonal antibodies that can bind to *Staphylococcus* of both coagulase types and that can enhance phagocytosis and killing of the bacteria and thereby enhance protection *in vivo*. There is also a need in the art for the epitope of the site to which such antibodies can bind so that other antibodies with similar abilities can be identified and isolated.

There is a related need in the art for humanized or other chimeric human/mouse monoclonal antibodies. In recent well publicized studies, patients administered murine anti-TNF (tumor necrosis factor) monoclonal antibodies developed anti-murine antibody responses to the administered antibody. (Exley A.R., et al., Lancet 335:1275-1277 (1990)). This type of immune response to the treatment regimen, commonly referred to as the HAMA response, decreases the effectiveness of the treatment and may even render the treatment completely ineffective. Humanized or chimeric human/mouse monoclonal antibodies have been shown to significantly decrease the HAMA response and to increase the therapeutic effectiveness. See LoBuglio et al., P.N.A.S. 86:4220-4224 (June 1989).

SUMMARY OF THE INVENTION

To address these needs in the art, the present invention encompasses opsonic and protective monoclonal and chimeric antibodies that bind to lipoteichoic acid of Gram positive bacteria. The antibodies also bind to whole bacteria and enhance phagocytosis and killing of the bacteria *in vitro* and enhance protection from lethal infection *in vivo*. The mouse monoclonal antibody has been humanized and the resulting chimeric antibody provides a previously unknown means to diagnose, prevent and/or treat infections caused by gram positive bacteria bearing lipoteichoic acids. This invention also encompasses a peptide mimic of the lipoteichoic acid epitope binding site defined by the monoclonal antibody. This epitope or epitope peptide mimic identifies other antibodies that may bind to the lipoteichoic acid epitope. Moreover, the epitope or epitope peptide mimic provides a valuable substrate for the generation of vaccines or other therapeutics.

BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 provides a schematic representation of lipoteichoic acid (LTA) in the Gram positive bacterial cell wall.

Figure 2 depicts antibody regions, such as the heavy chain constant region (C_H), the heavy chain variable region (V_H), the light chain constant region (C_L), and the light chain variable region (V_L).

Figure 3 demonstrates the enhancement of survival after administration of MAB 96-110 in a lethal neonatal model of coagulase positive staphylococcus sepsis.

Figure 4 demonstrates enhancement of survival in adult mice infected with coagulase negative staphylococci after administration of MAB 96-110. Approximately 23 hours after infection, 70% of the animals treated with MAB 96-110 were alive compared with 20% of animals not given antibody.

Figure 5 (SEQ ID NOS 4 & 5, and 6 & 7, respectively) provides a list of 18 resulting sequences for the 6mer library panning.

Figure 6 (SEQ ID NOS 8-43, respectively) provides a list of the 18 resulting sequences for the second experiment 15mer library panning.

Figure 7 (SEQ ID NOS 44 & 45, and 46 & 47, and 48 & 49, respectively) provides a list of the 17 resulting sequences for the first experiment 15mer library panning.

Figure 8 (SEQ ID NOS 50-67, respectively) provides a master list compiled from the common resulting peptide sequences from all the pannings.

Figure 9 sets forth a comparison of the optical density signals of each phage isolate at 6.25×10^{11} vir/ml.

Figure 10 shows the strategy for cloning the variable region gene fragments.

Figure 11 (SEQ ID NOS 68-85, respectively) lists the oligonucleotides primers used.

Figure 12 (SEQ ID NOS 86-105, respectively) provides the final consensus DNA sequence of the heavy and light chain variable regions.

Figure 13 demonstrates the re-amplification of the variable region gene fragments.

Figure 14 sets forth the plasmid map for pJRS334.

Figure 15 provides the results of the antibody production assay, showing that the transfection of cells with the plasmid construct results in the production of a molecule containing both human IgG and kappa domains.

Figure 16 provides the results of the activity assay, demonstrating that the transfection of cells with the plasmid construct results in the production of a molecule that binds to the Hay antigen.

Figure 17 depicts the opsonic activity of the chimeric monoclonal antibody 96-110 for *S. epidermidis* in a neutrophil mediated opsonophagocytic bactericidal assay.

Figure 18 demonstrates the enhancement of survival after administration of MAB 96-110 in a lethal model of *S. epidermidis* sepsis.

Figure 19 depicts the effect of the chimeric monoclonal antibody 96-110 on the survival of adult mice after intraperitoneal challenge with *S. epidermidis*.

Figure 20 sets forth the effect of the chimeric monoclonal antibody 96-110 on bacteremia in a lethal *S. epidermidis* model.

Figure 21 depicts bacteremia levels four hours after infection with *S. epidermidis* at different doses of the chimeric monoclonal antibody 96-110.

Figure 22 sets forth the effect of the chimeric monoclonal antibody 96-110 on survival in a lethal neonatal *S. epidermidis* sepsis model.

Sensitivity analysis	
Model 1	0.00
Model 2	0.00
Model 3	0.00
Model 4	0.00
Model 5	0.00
Model 6	0.00
Model 7	0.00
Model 8	0.00
Model 9	0.00
Model 10	0.00
Model 11	0.00
Model 12	0.00
Model 13	0.00
Model 14	0.00
Model 15	0.00
Model 16	0.00
Model 17	0.00
Model 18	0.00
Model 19	0.00
Model 20	0.00
Model 21	0.00
Model 22	0.00
Model 23	0.00
Model 24	0.00
Model 25	0.00
Model 26	0.00
Model 27	0.00
Model 28	0.00
Model 29	0.00
Model 30	0.00
Model 31	0.00
Model 32	0.00
Model 33	0.00
Model 34	0.00
Model 35	0.00
Model 36	0.00
Model 37	0.00
Model 38	0.00
Model 39	0.00
Model 40	0.00
Model 41	0.00
Model 42	0.00
Model 43	0.00
Model 44	0.00
Model 45	0.00
Model 46	0.00
Model 47	0.00
Model 48	0.00
Model 49	0.00
Model 50	0.00
Model 51	0.00
Model 52	0.00
Model 53	0.00
Model 54	0.00
Model 55	0.00
Model 56	0.00
Model 57	0.00
Model 58	0.00
Model 59	0.00
Model 60	0.00
Model 61	0.00
Model 62	0.00
Model 63	0.00
Model 64	0.00
Model 65	0.00
Model 66	0.00
Model 67	0.00
Model 68	0.00
Model 69	0.00
Model 70	0.00
Model 71	0.00
Model 72	0.00
Model 73	0.00
Model 74	0.00
Model 75	0.00
Model 76	0.00
Model 77	0.00
Model 78	0.00
Model 79	0.00
Model 80	0.00
Model 81	0.00
Model 82	0.00
Model 83	0.00
Model 84	0.00
Model 85	0.00
Model 86	0.00
Model 87	0.00
Model 88	0.00
Model 89	0.00
Model 90	0.00
Model 91	0.00
Model 92	0.00
Model 93	0.00
Model 94	0.00
Model 95	0.00
Model 96	0.00
Model 97	0.00
Model 98	0.00
Model 99	0.00
Model 100	0.00

The antibodies of the invention are broadly reactive with Gram positive bacteria, meaning that they selectively recognize and bind to Gram positive bacteria and do not recognize or bind to Gram negative bacteria. Any conventional binding assay can be used to assess this binding, including for example, the enzyme linked immunosorbent assay described below. The basis of the binding is the presence of LTA exposed on the surface of the cell wall of Gram positive bacteria.

10

species, in general teichoic acids make up a major part of the cell wall. The teichoic acids related to this invention are lipoteichoic acids which are teichoic acids made up of glycerol phosphate which is primarily linked to a glycolipid in the underlying cell membrane. Although the precise structure of LTA in the Gram positive bacterial cell wall is not known, a standard schematic representation commonly accepted in the art is set forth in Figure 1. Accordingly, the antibodies of the claimed invention are broadly reactive because they recognize and bind to the lipoteichoic acids that are characteristically surface exposed on Gram positive bacteria.

The antibodies of the invention are also opsonic, or exhibit opsonic activity, for Gram positive bacteria. As those in the art recognize, "opsonic activity" refers to the ability of an opsonin (generally either an antibody or the serum factor C3b) to bind to an antigen to promote attachment of the antigen to the phagocyte and thereby enhance phagocytosis. Certain bacteria, especially encapsulated bacteria which resist phagocytosis due to the presence of the capsule, become extremely attractive to phagocytes such as neutrophils and macrophages when coated with an opsonic antibody and their rate of clearance from the bloodstream is strikingly enhanced. Opsonic activity may be measured in any conventional manner as described below.

The ability of the anti-LTA antibodies of the invention to bind to and opsonize Gram positive bacteria and thereby enhance phagocytosis and cell killing *in vitro* and to enhance protection *in vivo* is completely unexpected because anti-LTA antibodies have been reported to lack opsonic activity. Indeed, anti-LTA antibodies have been often used as controls.

For example, Fattom et al., J. Clin. Micro. 30(12):3270-3273 (1992) examined the opsonic activity of antibodies induced against type specific capsular polysaccharide of *S. epidermidis*, using as controls antibodies induced against teichoic acids and against *S. hominus*. While type-specific antibodies were highly opsonic, anti-teichoic acid antibodies were not different from the anti-*S. hominus* antibodies.

Similarly, in Kojima et al., J. Infect. Dis. 162:435-441 (1990), the authors assessed the protective effects of antibody to capsular polysaccharide/adhesion against catheter-related bacteremia due to coagulase negative Staphylococci and specifically used a strain of *S. epidermidis* that expresses teichoic acid as a control. See page 436, Materials and Methods, left column, first ¶; right column, third ¶. In a later study, the authors reached a more explicit conclusion against the utility of anti-teichoic antibodies:

Immunization protocols designed to elicit antibody to teichoic acid but not to PS/A afforded no protection against bacteremia or endocarditis.

Takeda, et al., Circulation 86(6):2539-2546 (1991).

Contrary to the prevailing view in the field, the broadly reactive opsonic antibodies against the LTA of Gram positive bacteria, including *S. aureus* and *S. epidermidis*, of the invention satisfy a clear need in the art. As described in the background section, both *S. aureus* and *S. epidermidis* are common causes of post-operative wound infections; *S. epidermidis* has become a major cause of nosocomial infections in patients whose treatments include the placement of foreign objects; *S. epidermidis* has become a common cause of peritonitis in patients with continuous

serotypes of *S. aureus* (Tables 3-6) but not to the Gram negative control, *Haemophilus influenza*.

Similar to the antibodies described in the Fischer applications and patent set forth above, the antibody of the present invention exhibits very strong binding, i.e., O.D.s of around twice background in an enzyme-linked immunosorbent assay (described below), against strain Hay. In a preferred embodiment, the level of high binding is equal to or greater than five times background. In other embodiments, the level of high binding is equal to or greater than 10 times background. Of course, any meaningful increase over background (the level observed when all the reagents other than the antibody being tested) will be recognized by skilled persons in the art as high binding and therefor within the scope of the invention.

Also as described in the Fischer applications and patent, high binding has been found to correlate with opsonic activity. As set forth in Example 2, in a neutrophil mediated bactericidal assay (described below), the 96-110 MAB exhibited enhanced opsonization against the prototypic coagulase negative bacteria, *S. epidermidis*, and against the prototypic coagulase positive bacteria, *S. aureus*. With this level of opsonic activity, an antibody should enhance phagocytosis and cell killing of both coagulase negative and coagulase positive bacteria.

The term "enhanced" refers to activity that measurably exceeds background at a valuable level. The level deemed valuable may well vary depending on the specific circumstances of the infection, including the type of bacteria and the severity of the infection. For example, for enhanced opsonic or phagocytic activity, in a preferred

chains that have specified three dimensional structural and specific charge characteristics.

In a series of panning experiments set forth in Examples 4-6, we identified peptide sequences to which MAB 96-110 bound strongly. These sequences provide at least peptide mimics of the epitope to which MAB 96-110 bound. Thus, one aspect of the present invention involves a peptide having the sequence

W R M Y F S H R H A H L R S P (SEQ ID NO 1)

and another aspect of the invention involves a peptide having the sequence

W H W R H R I P L Q L A A G R (SEQ ID NO 2).

Of course, the epitope of the invention may be identical to one of these sequences or may be substantially homologous to these sequences such that the anti-LTA antibodies of the invention will bind to them. Alternatively, the substantially homologous sequences of the invention are those that are able to induce the anti-LTA antibodies of the invention. Other peptide epitope mimics within the invention may vary in length and sequence from these two peptides.

The present invention also encompasses recombinant epitopes, epitope mimics, and antigens. The DNA sequence of the gene coding for the isolated antigen can be identified, isolated, cloned, and transferred to a prokaryotic or eukaryotic cell for expression by procedures well-known in the art. For example, procedures are generally described in Sambrook et al., Molecular Cloning, A Laboratory Manual, 2nd Ed., Cold Springs Harbor Press, Cold Spring Harbor, New York (1989) incorporated by reference.

derivatives of the present invention are capable of enhancing the opsonization of such bacteria, at rates ranging from 75% and up.

The "fragments" of the antibodies of the invention include, for example, Fab, Fab', F(ab')₂, and SFv. These fragments are produced from intact antibodies using methods well known in the art such as, for example, proteolytic cleavage with enzymes such as papain (to produce Fab fragments) or pepsin (to produce F(ab')₂). The "regions" of the antibodies of the present invention include a heavy chain constant region (H_c or C_H), a heavy chain variable region (H_v or V_H), a light chain constant region (L_c or C_L), and a light chain variable region (L_v or V_L) (Figure 2). The light chains may be either a lambda or a kappa chain.

In a preferred aspect of the invention, the regions include at least one heavy chain variable region or a light chain variable region which binds a portion of LTA, including for example the specific antigen binding sites (that which binds to the epitope) of the two regions. In another embodiment, these two variable regions can be linked together as a single chain antibody. While a full length heavy chain may be critical for opsonic activity and enhance anti-cytokine (anti-inflammatory) activity, the antibody fragments encompassing the variable regions may be suitable for inhibition of bacterial binding to epithelial cells and may also be anti-inflammatory.

In a particularly preferred aspect of the invention, the antibody is a chimeric mouse/human antibody made up of regions from the anti-LTA antibodies of the invention together with regions of human antibodies. For example, a chimeric H chain can comprise the antigen binding region of the heavy chain variable region of the anti-

LTA antibody of the invention linked to at least a portion of a human heavy chain constant region. This humanized or chimeric heavy chain may be combined with a chimeric L chain that comprises the antigen binding region of the light chain variable region of the anti-LTA antibody linked to at least a portion of the human light chain constant region.

The chimeric antibodies of the invention may be monovalent, divalent, or polyvalent immunoglobulins. For example, a monovalent chimeric antibody is a dimer (HL) formed by a chimeric H chain associated through disulfide bridges with a chimeric L chain, as noted above. A divalent chimeric antibody is a tetramer ($H_2 L_2$) formed by two HL dimers associated through at least one disulfide bridge. A polyvalent chimeric antibody is based on an aggregation of chains.

A particularly preferred chimeric antibody of the invention is described in Examples 8-10 which set forth in detail the preparation of a preferred chimeric IgG antibody (and in Examples 11-13 which describe the functional activity of this preferred chimeric antibody). Of course, other chimeric antibodies composed of different sections of the anti-LTA antibodies of the invention are within the invention. In particular, the heavy chain constant region can be an IgM or IgA antibody.

In addition to the protein fragments and regions of the antibodies, the present invention also encompasses the DNA sequence of the gene coding for the antibodies as well as the peptides encoded by the DNA. Particularly preferred DNA and peptide sequences are set forth in Figure 12. That figure provides the variable regions of both the heavy and light chains of MAB 96-110, including the Complementarity Determining

The present invention also discloses a pharmaceutical composition comprising the anti-LTA antibodies, whether polyclonal, monoclonal or chimeric, as well as fragments, regions, and derivatives thereof, together with a pharmaceutically acceptable carrier. The pharmaceutical compositions of the invention may alternatively comprise the isolated antigen, epitope, or portions thereof, together with a pharmaceutically acceptable carrier.

Pharmaceutically acceptable carriers can be sterile liquids, such as water, oils, including petroleum oil, animal oil, vegetable oil, peanut oil, soybean oil, mineral oil, sesame oil, and the like. With intravenous administration, water is a preferred carrier. Saline solutions, aqueous dextrose, and glycerol solutions can also be employed as liquid carriers, particularly for injectable solutions. Suitable pharmaceutical carriers are described in Remington's Pharmaceutical Sciences, 18th Edition (A. Gennaro, ed., Mack Pub., Easton, Pa., 1990), incorporated by reference.

Finally, the present invention provides methods for treating a patient infected with, or suspected of being infected with, a Gram positive bacteria such as a staphylococcal organism. The method comprises administering a therapeutically effective amount of a pharmaceutical composition comprising the anti-LTA immunoglobulin (whether polyclonal or monoclonal or chimeric, including fragments, regions, and derivatives thereof) and a pharmaceutically acceptable carrier. A patient can be a human or other mammal, such as a dog, cat, cow, sheep, pig, or goat. The patient is preferably a human.

Yet another preferred embodiment of the present invention is a vaccine comprising the epitope, epitope mimic, or other part of the LTA antigen and a pharmaceutically acceptable carrier. Upon introduction into a host, the vaccine generates an antibody broadly protective and opsonic against infection by Gram positive bacteria. The vaccine may include the epitope, an epitope mimic, any mixture of epitopes and epitope mimics, the antigen, different antigens, or any combination of epitopes, epitope mimics and antigens.

Vaccinations are particularly beneficial for individuals known to be or suspected of being at risk of infection by Gram positive bacteria. This includes patients receiving body implants, such as valves, patients with indwelling catheters, patients preparing to undergo surgery involving breakage or damage of skin or mucosal tissue, certain health care workers, and patients expected to develop impaired immune systems from some form of therapy, such as chemotherapy or radiation therapy.

Treatment comprises administering the pharmaceutical composition (including antibodies and vaccines) by intravenous, intraperitoneal, intracorporeal injection, intra-articular, intraventricular, intrathecal, intramuscular, subcutaneous, intranasally, intravaginally, orally, or by any other effective method of administration. The composition may also be given locally, such as by injection to the particular area infected, either intramuscularly or subcutaneously. Administration can comprise administering the pharmaceutical composition by swabbing, immersing, soaking, or wiping directly to a patient. The treatment can also be applied to objects to be placed within a patient, such as dwelling catheters, cardiac valves, cerebrospinal fluid shunts,

joint prostheses, other implants into the body, or any other objects, instruments, or appliances at risk of becoming infected with a Gram positive bacteria, or at risk of introducing such an infection into a patient.

As a particularly valuable corollary of treatment with the compositions of the invention (including all anti-LTA antibodies (whether polyclonal or monoclonal or chimeric, including fragments, regions, and derivatives thereof), all pharmaceutical compositions based on such antibodies, as well as on epitope, epitope mimics, or other part of the LTA antigen and vaccines based on such epitope or antigens) is the reduction in cytokine release that results from the introduction of the LTA of a Gram positive bacteria. As is now recognized in the art, LTA induces cytokines, including for example tumor necrosis factor alpha, Interleukin 6, and interferon gamma. See Takada et al., *Infection and Immunity*, 63 (1):57-65 (January 1995). Accordingly, the compositions of the invention may enhance protection at three levels: (1) by binding to LTA on the bacteria and thereby blocking the initial binding to epithelial cells and preventing subsequent invasion of the bacteria; (2) by enhancing opsonization of the bacteria and thereby enhancing clearance of the bacteria from tissues and blood; and/or (3) by binding to LTA and partially or fully blocking cytokine release and modulating the inflammatory responses to prevent shock and tissue destruction.

Having generally described the invention, it is clear that the invention overcomes some of the potentially serious problems described in the Background regarding the development of antibiotic resistant Gram positive bacteria. As set forth above, Staphylococci and streptococci (such as *S. faecalis*) have become increasingly

(MeOH). One hundred microliters of the suspension in MeOH was distributed into each well of Nunc Maxisorp Stripwells. The MeOH was allowed to evaporate, fixing the bacteria to the plastic. The bacteria-coated stripwells were stored in plastic bags and used within 2 months of preparation.

For evaluation of antibodies, the bacteria-coated plates were washed once with PBS and non-specific reactive sites on the bacteria were blocked by the addition of 120 ul/well of a solution of 1% bovine serum albumin (BSA) in PBS. After a 30-60 minute incubation, the wells were washed four times with PBS containing 0.05% Tween-20 (PBS-T). Antibody, diluted in PBS-T, was then added to the wells. Supernatants were tested at a dilution of 1:2. Ascites and purified antibody were tested at dilutions indicated in the Tables. After addition of the antibody, the wells were incubated at room temperature for 30-60 minutes in a draft-free environment. The wells were again washed four times with PBS-T and each well received 95 ul of detection antibody: rabbit anti-mouse IgG, conjugated to horse radish peroxidase (HRP) and diluted 1:4000 in PBS-T. The detection antibodies were specific for mouse gamma, mu or alpha heavy chains (Zymed catalog numbers 61-6020, 61-6820 or 61-6720 respectively), as indicated in the Tables.

Following another 30-60 incubation at room temperature, the wells were washed four times with PBS-T and each well received 100 ul of one-component TMB substrate solution (Kirkegaard and Perry Labs catalog number 50-76-05). The wells were incubated in the dark at room temperature for 15 minutes. The reaction was stopped by the addition of 80 ul of TMB stop solution (Kirkegaard and Perry Labs catalog

number 50-85-05) and the absorbance of each well at 450 nm was determined using a Molecular Devices Vmax plate reader.

Immunoassay on LTA's: Reactivity of MAB 96-110 was measured by immunoassay on wells coated with LTA's. LTA's were obtained from Sigma Chemical Company and diluted in PBS to 1 ug/ml. One hundred microliters of the 1 ug/ml solution was distributed into replicate Nunc Maxisorp Stripwells. The LTA was incubated in the wells overnight at room temperature. The unbound material was removed from the wells with four washes of PBS-T. The wells were not blocked with BSA or other proteins. Antibody, diluted in PBS-T, was then added to the wells and the assay continued as described above.

Competitive Inhibition of Antibody of LTA: In order to determine the ability of LTA to inhibit binding of MAB 96-110 to wells coated with MeOH-fixed Strain Hay, a competitive inhibition assay was performed. Wells were coated in MeOH with Strain Hay and blocked with BSA as described above. Fifty ul of LTA from *S. mutans*, *S. aureus* or *S. faecalis* were added to duplicate wells. Six different concentrations of each LTA were tested (from 0.04 to 9.0 ug/ml). LTA's were diluted in PBS-T to obtain the desired concentrations. Immediately after addition of the LTA, 50 ul of purified MAB 96-110 at 1 ug/ml was added to each well. The final dilution of the MAB 96-110 was therefore 0.5 ug/ml. Uninhibited control wells received only PBS-T and MAB without LTA.

Binding of MAB 96-110 to the LTA in the PBS-T solution resulted in a complex of MAB/LTA which was removed from the plate during the subsequent washing step. The

interaction of the MAB 96-110 with the LTA inhibited the antibody from binding to the LTA on the surface of the bacteria and thus reduced the number of MAB 96-110 molecules bound to the MeOH-fixed strain Hay used to coat the wells. Because the number of MAB 96-110 molecules bound to the MeOH-fixed Strain Hay was reduced, the level of binding of the detection antibody (rabbit anti-mouse IgG-HRP) was therefore also decreased, leading to a reduction of color development when compared to wells in which no LTA was present.

Immunoassay with Protein A Method: In order to evaluate monoclonal antibody 96-110 for reactivity with *S. aureus* 5 and *S. aureus* 8, it was necessary to modify the immunoassay procedure described above. Both *S. aureus* strains express Protein A on their surfaces. Because Protein A binds strongly to the constant region of the heavy chains of gamma-globulins, it was possible that false positive results would be obtained due to non-specific binding of the 96-110 antibody to the Protein A molecule. In order to overcome this difficulty, the immunoassay wells were coated with bacteria as described above. However, prior to the addition of the 96-110 antibody to the bacteria-coated wells, the purified monoclonal antibody (MAb) was reacted with a solution of recombinant Protein A conjugated to HRP and diluted 1:500 in PBS-T. This reaction was allowed to proceed for 30 minutes. The wells were washed four times with PBS-T and 100 ul of the solution of Protein A-HRP-MAb was added to the wells. The presence of the Protein A-HRP from the pretreatment prevented the MAb from binding to the Protein A on the *S. aureus* 5 and 8. Furthermore, the binding of the Protein A-HRP to the constant region of the heavy chain did not interfere with the antibody

catheter, to more closely mimic the clinical setting. An alternative model utilizes adult susceptible animals, such as CF1 mice.

Clearance is evaluated by determining whether the pharmaceutical composition enhances clearance of the infectious agent from the animal. This is typically determined from a sample of biological fluid, such as blood, peritoneal fluid, or cerebrospinal fluid. The infectious agent is cultured from the biological fluid in a manner suitable for growth or identification of the surviving infectious agent. From samples of fluid taken over a period of time after treatment, one skilled in the art can determine the effect of the pharmaceutical composition on the ability of the animal to clear the infectious agent. Further data may be obtained by measuring over a period of time, preferably a period of days, survival of animals to which the pharmaceutical composition is administered. Typically, both sets of data are utilized. Results are considered positive if the pharmaceutical composition enhances clearance or decreases mortality. In situations in which there is enhanced organism clearance, but the test animals still perish, a positive result is still indicated.

EXAMPLE 1

The Production of Hybridomas and Monoclonal Antibodies

To produce monoclonal antibodies that were directed against the surface proteins of *S. epidermidis* and were opsonic and protective for *S. epidermidis*, mice were immunized with whole *S. epidermidis*, Strain Hay.

A suspension of heat killed *S. epidermidis* was adjusted to an optical density (OD) of 0.137 at a wavelength of 650 nm when measured through a 1 centimeter light path. Bacteria from five mls of the suspension were pelleted by centrifugation (approximately 1800 X g, 10 minutes, room temperature). The supernatant was discarded and the pellet resuspended in 0.6 mls of PBS, which was then mixed with 0.9 mls of complete Freund's adjuvant (CFA). The resulting emulsion was used as the immunogen.

Adult, female BALB/c mice, obtained from Harlan Sprague Dawley (Indianapolis, IN) were immunized subcutaneously with 0.2 mls of the immunogen described above. The mice received a booster immunization approximately two and ½ months later with antigen prepared as described above, except that incomplete Freund's adjuvant (IFA) was used as the adjuvant instead of CFA. A final, prefusion boost was given approximately two months after that. This boost consisted of 1 ml of *S. epidermidis* suspension ($OD_{650}=0.137$). Mice 8159 and 8160 each received an intraperitoneal injection of 0.5 mls of the suspension. Five days later, the spleen from mouse 8159 was removed and used for hybridoma formation.

Hybridomas were prepared by the general methods of Shulman, Wilde and Kohler Nature 276:269-270 (1978) and Bartal and Hirshaut "Current Methods in Hybridoma Formation in Methods of Hybridoma Formation, Bartal and Heishaut, eds., Humana Press, Clifton, New Jersey (1987). A total of 2.135×10^8 spleenocytes from mouse 8159 were mixed with 2.35×10^7 SP2/0 mouse myeloma cells (ATCC Catalog number CRL1581) and pelleted by centrifugation (400 X g, 10 minutes at room temperature) and washed in serum free medium. The supernatant was removed to near-dryness and fusion of the cell mixture was accomplished in a sterile 50 ml centrifuge conical by the addition of 1 ml of polyethylene glycol (PEG; mw 1400; Boehringer Mannheim) over a period of 60-90 seconds. The PEG was diluted by slow addition of serum-free medium in successive volumes of 1, 2, 4, 8, 16 and 19 mls. The hybridoma cell suspension was gently resuspended into the medium and the cells pelleted by centrifugation (500 X g, 10 minutes at room temperature). The supernatant was removed and the cells resuspended in medium RPMI 1640, supplemented with 10% heat-inactivated fetal bovine serum, 0.05 mM hypoxanthine and 16 uM thymidine (HT medium). One hundred ul of the hybridoma cells were planted into 760 wells of 96-well tissue culture plates. Eight wells (column 1 of plate A) received approximately 2.5×10^4 SP2/0 cells in 100 ul. The SP2/0 cells served as a control for killing by the selection medium added 24 hours later.

Twenty four hours after preparation of the hybridomas, 100 ul of RPMI 1640, supplemented with 10% heat-inactivated fetal bovine serums, 0.1 mM hypoxanthine, 0.8 uM aminopterin and 32 uM thymidine (HAT medium) was added to each well.

Ninety six hours after the preparation of the hybridomas, the SP2/0 cells in plate A, column 1 appeared to be dead, indicating that the HAT selection medium had successfully killed the unfused SP2/0 cells.

Eleven days after the preparation of the hybridomas, supernatants from all wells were tested by ELISA for the presence of antibodies reactive with methanol-fixed *S. epidermidis*. Based on the results of this preliminary assay, cells from 20 wells were transferred to a 24-well culture dish. Four days later, supernatant from these cultures were retested by ELISA for the presence of antibodies reactive with methanol-fixed *S. epidermidis*. Of these supernatants, one (from colony 96-105CE11) was a strongly reactive IgG (Table 1). Two colonies (96-105FD4 and 96-105GB5) were very weakly reactive IgG's and one colony 96-105HB10 was a weakly reactive IgM. Antibodies of the IgM isotype are not as desirable as IgG's and culture 96-105HB10 was cryopreserved and not further examined.

Cultures 96-105 CE11, FD4 and GB5 were reanalyzed several days later and only CE11 showed a strong response (Table 2). No response was obtained with the other cell cultures, and no further experimental work was done with them.

To further test the specificity of this antibody for *S. epidermidis*, a whole cell ELISA with several bacteria was performed (Table 3). The antibodies from this colony bound strongly to *S. epidermidis* (Hay) O.D. 1.090 and to a lesser degree to Group B streptococcus (GBS), but not to *H. influenzae* (HIB+, with type b capsule; HIB- without typable capsule) or type 14, pneumococcus (Pn 14).

A clone from 96-105CE11 IF6 was isolated and retested and was an IgG-1 that reacted strongly with *S. epidermidis* (Strain Hay) in the whole cell ELISA (Table 4). This clone was then designated 96-110. To determine if 96-110 had the broad binding characteristics we sought and would be consistent with binding to the surface protein on *S. epidermidis* (Strain Hay) that bound broadly opsonic antibody, we ran a whole cell ELISA against several coagulase negative staphylococci (Table 5). Using 96-110 in Ascites fluid, strong binding at several dilutions was detected for *S. epidermidis* type I, II, III, *S. hemolyticus* and *S. hominus*.

In addition, 96-110 MAB was purified over a protein G column (Pharmacia). Using a modification of the whole cell ELISA, peroxidase labeled protein A was mixed with the purified 96-110 MAB and then reacted with *S. aureus* type 5 (SA5) and *S. aureus* type 8 (SA8) obtained from ATCC at Accession Nos. 12602 and 12605, respectively. Both *S. aureus* serotypes reacted strongly with the 96-110 MAB (Table 6). Since, in our previous studies, we found that absorption with *S. epidermidis* (Strain Hay) could decrease IgG opsonic activity and opsonic antibodies raised against Hay reacted with a surface protein of Hay, we felt that this was still consistent with a MAB to the surface protein we were trying to characterize. This finding was also important since types 5 & 8, *S. aureus* are serotypes commonly associated with human infections. Using this protein A assay, MAB to type 14 pneumococcus did not demonstrate binding to *S. aureus*.

Therefore, we have identified a mouse IgG₁ MAB raised against *S. epidermidis* (Strain Hay) that binds to the surface of both coagulase negative and coagulase

TABLE 1

Immunoassay Results, 96-105 Supernatants on
Methanol-Fixed S. Hay

Colony ID	Detection Specific For:		
	G	A	M
PBS-F	0.070	0.080	0.050
CE11	0.788	0.065	0.056
EB5	0.079	0.065	0.053
EE5	0.084	0.069	0.055
FD4	0.089	0.067	0.059
FG4	0.087	0.065	0.065
FG8	0.090	0.060	0.062
FF9	0.095	0.062	0.059
GE4	0.074	0.067	0.059
GB5	0.155	0.077	0.078
GB6	0.073	0.062	0.053
GC6	0.069	0.062	0.052
GC9	0.076	0.062	0.053
GB10	0.075	0.064	0.102
HG2	0.195	0.067	0.059
HG3	0.079	0.066	0.060
HE4	0.076	0.073	0.065
HG4	0.077	0.101	0.061
HG5	0.077	0.062	0.058
HC8	0.083	0.064	0.057
HB10	0.070	0.064	0.223
AC4 IID10*	0.065	0.066	0.069

*Monoclonal antibody reactive with Hib protein D.

Immunoassay Results, 96-105 Supernatants on
Methanol-Fixed S. Hay

Colony ID	Detection Specific For:		
	G	A	M
Buffer	0.052	0.045	0.045
CE11	0.933	0.049	0.046
FD4	0.073	0.054	0.051
GB5	0.050	0.040	0.036

Variable	Mean	SD	Min	Max
Age	34.5	10.2	22	55
Gender	0.5	0.5	0	1
Marital status	0.6	0.5	0	1
Education	12.5	1.5	10	16
Income	15.2	5.8	10	25
Health status	0.7	0.4	0	1
Employment	0.8	0.4	0	1
Family size	3.2	1.8	1	8
Home ownership	0.9	0.3	0	1
Auto ownership	0.7	0.4	0	1
Life satisfaction	4.2	1.5	1	7
Health satisfaction	5.1	1.2	1	7
Income satisfaction	3.8	1.8	1	7
Employment satisfaction	4.5	1.4	1	7
Family size satisfaction	3.5	1.6	1	7
Home ownership satisfaction	5.5	1.1	1	7
Auto ownership satisfaction	4.8	1.3	1	7
Life satisfaction (control)	4.3	1.4	1	7
Health satisfaction (control)	5.2	1.1	1	7
Income satisfaction (control)	3.9	1.7	1	7
Employment satisfaction (control)	4.6	1.3	1	7
Family size satisfaction (control)	3.6	1.5	1	7
Home ownership satisfaction (control)	5.6	1.0	1	7
Auto ownership satisfaction (control)	4.9	1.2	1	7

TABLE 3

Immunoassay Results, 96-105 Supernatants on
Methanol-Fixed Bacteria

Colony ID	Detection Antibody	Hay	Hib+	Hib-	GBS	Pn14
CE11	gamma-specific	1.090	0.106	0.068	0.304	0.063
FE11	gamma-specific	0.167	0.084	0.068	0.112	0.053
Buffer	gamma-specific	0.048	0.075	0.056	0.070	0.053

Several colonies from 96-105 not cloned.

1996-1997 "AT 94-6860

TABLE 4
Assay of 96-105 CE11 IF6 on Various Bacteria

Antibody	Antigen	Dilution	Isotype	Hay	Pn14
PBS-T				0.072	0.064
96-105CE11-IF6	Hay	2	IgG-1, k	1.608	0.099
		4		1.184	0.087
		8		0.846	0.069
		16		0.466	0.074

TABLE 5

Detection of Bacteria of Anti-Hay Monoclonal*
in Whole Cell ELISA

Antibody	Dilution	Staph. Hay	Staph. Epi I	Staph. Epi II	Staph. Epi III	Staph. Hemmolyt.	Staph. Hominus
Buffer		0.056	0.063	0.066	0.055	0.058	0.074
96-110 Ascites	100	1.448	2.334	1.524	1.241	1.197	0.868
	400	1.325	2.542	0.746	0.425	0.830	0.422
	1600	1.087	2.452	0.369	0.176	0.680	0.185
	6400	0.930	2.430	0.195	0.089	0.602	0.110
	25600	0.674	1.672	0.113	0.069	0.647	0.081

*Anti-Hay Monoclonal from unpurified ascites fluid

TABLE 6

Detection of Methanol-Fixed SA5, SA8 and S. Hay
By Purified Monoclonal Anti-Hay Using Protein A

Anti-Hay Dilution	ATCC SA5	ATCC SA8	USU Hay
500	1.329	3.345	3.017
1000	1.275	2.141	2.266
2000	0.873	1.016	1.487
4000	0.333	0.491	0.951
8000	0.159	0.232	0.490
16000	0.132	0.149	0.331
Normal Mouse 1000	0.101	0.090	0.082
Buffer	0.102	0.113	0.152

Purified anti-Hay stock = 1.63 mg/ml

EXAMPLE 2

The Opsonic Activity of the Monoclonal Antibody

Antibodies which bind to an antigen may not necessarily enhance opsonization or enhance protection from infection. Therefore, a neutrophil mediated bactericidal assay was used to determine the functional activity of antibody to *S. epidermidis*. Neutrophils were isolated from adult venous blood by dextran sedimentation and ficoll-hypaque density centrifugation. Washed neutrophils were added to round-bottomed wells of microtiter plates (approximately 10^6 cells per well) with approximately 3×10^4 mid-log phase bacteria (*S. epidermidis* Hay, ATCC 55133). Newborn lamb serum (10 uls), screened to assure absence of antibody to *S. epidermidis*, was used as a source of active complement.

Forty microliters of immunoglobulin (or serum) were added at various dilutions and the plates were incubated at 37°C with constant, vigorous shaking. Samples of 10 uls were taken from each well at zero time and after 2 hours of incubation. Each was diluted, vigorously vortexed to disperse the bacteria, and cultured on blood agar plates overnight at 37°C to quantitate the number of viable bacteria. Results are presented as percent reduction in numbers of bacterial colonies observed compared to control samples.

Since the 96-110 MAB bound to both coagulase negative and coagulase positive Staphylococci, opsonic studies were performed to determine if the MAB enhanced phagocytosis and killing of both groups of staphylococci. At a 1:80 dilution, the MAB

enhanced opsonization and killing of coagulase negative *Staphylococcus* (*S. epidermidis*) to 100%, compared with 49.5% with C' and PMN alone (Table 7). The coagulase positive *Staphylococcus* also showed enhanced phagocytosis at 1:10 and 1:40 dilution (83.3% and 78.9% respectively) compared with 53.7 percent with C' and PMN alone. At 1:80 dilution, the opsonic activity against *S. aureus* was decreased to 61%.

These data show that not only does the MAB bind to the surface of both coagulase negative and coagulase positive *Staphylococci*, but that it has functional activity and can enhance phagocytosis and killing of these bacteria. Such an antibody would be capable of promoting clearance of *Staphylococci* that have invaded a host and would be useful therapeutic agent.

TABLE 7

Opsonic Assay

Antibody: Purified M X Hay, 96-110

Group Description	Ab Dilution	% Killed <u><i>S. epidermidis</i></u>	% Killed <u><i>S. aureus</i></u>
C' only		0.0	0.0
PMN only		0.0	0.0
PMN + C' No Ab		49.5	53.7
PMN + Ab + C'	10	-	83.3
	40	-	78.9
	80	100.0	61.0

EXAMPLE 3

In vivo Protective Efficacy

Opsonic antibody correlates with enhanced protection from staphylococcal infections, as set forth in the recent series of Fischer applications and issued patent described and incorporated by reference above. To further demonstrate that the MAB can enhance survival to infections with both coagulase positive and coagulase negative Staphylococci, studies were conducted using lethal infection models.

Two day old Wistar rats were injected with 10^6 *S. aureus* (type 5, ATCC 12605) subcutaneously just cephalad to the tail. Approximately 30 minutes before and 24 and 48 hours after infection, 0.2 ml MAB 96-110 (~320 ug) was given IP. Control animals were given an equal volume of saline or a control MAB not directed against Staphylococci. All animals were observed daily for five days to determine survival.

MAB 96-110 enhanced survival in this lethal neonatal model of coagulase positive staphylococcus sepsis (Figure 3): 8/15 survived after treatment with MAB 96-110, and 0/10 survived with Control MAB or 2/25 with saline treatment.

In a similar manner MAB 96-110 enhanced survival in adult mice infected with coagulase negative staphylococci. Adult CF1 mice were given 0.5 ml *S. epidermidis* (Hay) IP (3.5×10^9 bacteria). About 24 hrs and 2 hrs before and 24 hrs post-infection, 320 ug of MAB 96-110 were given to one group of mice and compared with a second group infected in the same manner, but not treated with antibody. All animals were followed 5 days to determine survival. Approximately 23 hours after infection, 70% of

the animals treated with MAB 96-110 were alive compared with 20% of animals not given antibody (Figure 4). When the study was terminated 50% of the MAB animals remained alive compared to only 10% of controls.

Thus, MAB 96-110 could enhance survival in lethal coagulase positive and coagulase negative staphylococcal infections. This enhancement occurred in an adult model and an immunocompromised model (immature neonatal immune system).

EXAMPLE 4

Peptide selection

Panning random 6mer and 15mer fd-tet phage libraries

Amplified random 6mer and 15mer fd-tet phage libraries were panned against the 96-110 antibody to yield populations of 6 and 15 amino acid length peptides which cross react with the 96-110 antibody. The original libraries were acquired from George P. Smith, Division of Biological Sciences, University of Missouri, Columbia, MO. In order to be used for panning, the 96-110 antibody was crosslinked to Biotin using the Sulfo-NHS-biotin ester crosslinking kit following the manufacturers protocol (Pierce Chemical Co.).

For the first round of panning, 35mm polystyrene petri dishes (Costar) were coated with streptavidin by incubating the plates overnight at 4°C rocking with 100mM NaHCO₃ and 10ug streptavidin. Streptavidin was then discarded and plates were filled with blocking solution (0.1M NaHCO₃, 5mg/ml dialyzed BSA, 0.1ug/ml streptavidin) and incubated for 1hr at 4°C. The following protocol was then followed: Wash dishes six

times with TBS/Tween (50mM Tris-HCl pH 7.5, 150mM NaCl, 0.5% v/v Tween 20). Incubate dishes overnight at 4°C rocking with 400ul TBS/Tween containing 1 mg/ml dialyzed BSA and 10ug biotinylated 96-110 antibody. Add 4ul 10mM biotin and allow to incubate 1hr at 4°C rocking. Wash dishes six times as previously stated. Add 400ul TBS/Tween into each dish, add 4ul 10mM biotin and add approximately 5ul of either the 6mer or 15mer amplified fd-tet phage library (at 1×10^{14} vir/ml). Rock dishes 4hrs at 4°C. Pour out phage and wash ten times with TBS/Tween. Incubate plates at room temp with 400ul elution buffer (0.1N HCl, pH adjusted to 2.2 with glycine, 1mg/ml BSA) for 10 min with rocking. Remove eluate to a Centricon 30 (Amicon) concentrator and buffer exchange with TBS (50mM Tris-HCl pH 7.5, 150mM NaCl) and concentrate to a volume of about 100ul. Amplify eluate by mixing 100ul eluate with 100ul K91 terrific broth cells and allowing phage to infect cells for 10-30min. Pipette infection mixture into 20ml pre-warmed NZY medium (10g NZ amine A, 5g yeast extract, 5g NaCl, 1 liter water, adjust to pH 7.5, autoclave) containing 0.2 ug/ml tetracycline. Shake vigorously at 37°C for 30-60 min. Add 20ul of 20 mg/ml tetracycline stock to the flask. Remove a small sample for titering on plates and allow flask to shake vigorously overnight at 37°C. Calculate yield from biopanning using the number of colonies counted on the titering plates from amplification infection and the number of input phage at the beginning of panning. This number should amount to at least approximately $10^{-5}\%$. Centrifuge 20ml culture for 10min at 5,000 rpm, then for 10min at 10,000rpm; pour the doubly cleared supernatant into a fresh tube containing 3ml PEG/NaCl (16.7% PEG 8000, 3.3M NaCl). Mix well and allow to incubate overnight at 4°C. Centrifuge tube 15

min at 10,000rpm, discard supernatant and redissolve phage pellet in 1ml TBS. Collect resuspended phage into a 1.5ml eppendorf tube, clarify the suspension by centrifugation, and add 150ul PEG/NaCl. Allow to incubate on ice for 1 hr. Microfuge the tube 10min, discard supernatant, and redissolve phage in 200ul TBS.

The second and third round of panning are carried out the same way. The eluted, amplified phage (100ul) from the previous panning is preincubated with biotinylated 96-110 antibody (100nM for the second round; 0.1nM for the third round) overnight at 4°C. 400 ul TBS/Tween is added to the mixture and it is pipetted onto streptavidin coated plates (prepared as previously stated) and then incubated with rocking gentle at room temperature for 10 min. The plates are then washed, eluted, and amplified as previously stated. The input and resultant phage are titered to determine yield from biopanning.

EXAMPLE 5

Sequencing resulting phage populations
to identify consensus sequences

After the third round of panning, the titered infection plates are used to pick 20 single isolated colonies for each library. The colonies are grown overnight in 5ml NZY medium containing 40ug/ml tetracycline. Replicative form DNA is extracted from each culture using Qiaplasamid quick prep kit (Qiagen Inc.) following the manufacturer's protocol. Media supernatants are saved for phage stock to be used in Example 4. 2.5ul of each RF DNA sample is added to a reaction containing 3.5pmole CLC502

primer (5'-TGAATTTTCTGTATGAGGTTT-3') (SEQ ID NO 3), 8 UI Prizm sequencing mix (ABI Inc.), QS to 20ul with water and amplified following manufacturers protocol. Successful sequences are translated and aligned. 18 resulting sequences for the 6mer library panning are listed in Figure 5. 18 resulting sequences for the second experiment 15mer library panning are listed in Figure 6. 17 resulting sequences for the first experiment 15mer library panning are listed in Figure 7. A master list was compiled of the common resulting peptide sequences from all the pannings (Figure 8) with the frequency of occurrence listed to the right of each sequence. Consensus portions of the sequences are marked on the master list (Figure 8).

EXAMPLE 6

Phage EIA comparing 3rd round phage isolates crossreactivity to 96-110 antibody

The saved media phage stocks for each of the common resulting peptide sequences were amplified as previously stated. Amplified phage preparations were quantitated by Abs₂₆₉ and diluted to 1×10^{13} vir/ml and serially diluted 1.2 seven times. A 96-well polystyrene plate was coated with 2ug/ml streptavidin in 0.1M NaHCO₃ overnight at 4°C. Plates were emptied and blocked for 1hr at room temperature with phage blocking solution, 100ul/well. The following protocol was then followed. Wash wells three times with TBS/Tween. Incubate overnight at 4°C with 0.2ug/ml biotinylated 96110 in phage blocking solution, 100ul/well. Wash wells three times with TBS/Tween. Incubate overnight at 4°C with serially diluted phage, 100ul/well. Wash wells three

times with TBS/Tween. Incubate 1 hr at room temperature with 1:5000 goat polyclonal anti-phage-HRP. Wash wells three times with TBS/Tween. Develop with 100ul ABTS substrate (Kirkegaard Perry) for 10-15min and read absorbance (402nm) on spectrophotometer according to manufacturer's protocol. Optical density signals of each phage isolate at 6.25×10^{11} vir/ml are compared in Figure 9. The two isolates yielding the greatest signals are:

15mer 2.12 W R M Y F S H R H A H L R S P (SEQ ID NO:1)

15mer 2.1 W H W R H R I P L Q L A A G R (SEQ ID NO:2)

EXAMPLE 7

Antibodies Against Lipoteichoic Acid (LTA)

As set forth above, we identified two peptides that reacted with 96-110 MAB. However, after identifying the peptides, the sequences did not correspond to any known proteins. Thus we began to consider other possible antigen candidates. We were surprised to find that MAB 96-110 bound strongly to LTA from several gram positive bacteria such as *S. mutans*, *S. aureus* and *S. faecalis* (Table 8). In addition, in an ELISA, when the wells coated with *S. epidermidis* (Strain Hay) were reacted with MAB 96-110 inhibited by varying concentrations of LTA (from *S. mutans*, *S. aureus* and *S. faecalis*), reduction in MAB binding occurred (Table 9). The inhibition of MAB 96-110 binding was greatest at the highest concentration of LTA inhibitor (9 ug/ml for each LTA) and varied according to which bacterial LTA was used (52% inhibition with *S. mutans*, 40.6% with *S. aureus* and 38.2% with *S. faecalis*).

The MAB 96-110 was also analyzed for binding to LTA from *S. pyogenes* (group A streptococcus) and various group A streptococcal M types. The MAB showed strong binding to the LTA and also bound to the different M types with strongest binding to M1 and M3 (Table 10).

We were surprised to find an antibody that bound to LTA and enhanced opsonization for both coagulase positive and coagulase negative staphylococci *in vitro* and enhanced survival in lethal models of staphylococcal (coagulase negative and coagulase positive) sepsis, *in vivo*. This is particularly surprising because the bacteria in each model were injected systemically (SQ or IP) and by-passed the epithelial barriers (skin or mucous membranes) where LTA is thought to possibly act as an adherence factor for the bacteria to epithelial cells.

In addition, this strong anti-LTA reactivity will provide a method to block the binding of LTA bearing bacteria to epithelial cells and prevent colonization of important pathogens such as staphylococci, group A streptococci, *S. faecalis* (enterococci) and *S. mutans*. Since LTA induces proinflammatory cytokines such as TNF, IL-6 and Interferon gamma, MABs with strong anti-LTA binding will also have an anti-inflammatory action and modulate cytokine production secondary to LTA bearing bacteria. Anti-LTA antibodies or vaccines could be designed and produced to modulate cytokine production and inflammation in tissues and prevent the adverse effects of these proinflammatory cytokines.

TABLE 8

Reactivity of Anti-Hay MAB 96-110 on wells Coated with Several LTA's

Antibody ID	Concentration or Dilution	LTA from <i>S. mutans</i>	LTA from <i>S.</i> <i>aureus</i>	LTA from <i>S.</i> - <i>faecalis</i>
Buffer	-	0.145	0.172	0.140
Anti-Hay	0.9 ug/ml	3.899	3.253	3.153
MAB	0.3 ug/ml	3.523	2.824	2.769
96-110	0.1 ug/ml	2.023	2.421	2.133
	0.033 ug/ml	2.143	1.590	1.539
	0.011 ug/ml	1.396	0.998	0.832

TABLE 9

Inhibition of Anti-Hay MAB 96-110
with LTA From Different Gram Positive Bacteria

LTA Inhibitor (ug/ml)	LTA <i>S. mutans</i>	LTA <i>S. aureus</i>	LTA <i>S. faecalis</i>
9	0.298	0.360	0.140
3	0.449	0.434	0.496
1	0.549	0.538	0.545
0.37	0.558	0.526	0.549
0.12	0.509	0.735	0.582
0.04	0.574	0.614	0.671
0	0.621	0.607	0.648

NOTES:

1. Wells were coated with methanol-fixed Hay.
2. Wells were blocked with 1% BSA in PBS.
3. Monoclonal anti-Hay was used at a final concentration of 0.5 ug/ml and reacted with inhibitors at the concentrations indicated in the Table.
4. Detection was with a gamma-specific Rabbit anti-Mouse.
5. Substrate was TMB.

TABLE 10

Reactivity of MAB 96-110 on Whole Methanol Fixed Group A Strep

Antibody ID	Dilution or Conc.	GAS* Type 1 #12344	GAS Type 3 #21546	GAS Type 18 #12357	GAS Type 24 #10782	Response on pyogenes LTA
Buffer		0.511	0.161	0.234	0.148	0.075
Anti-Hay MAb	0.3 ug/ml	1.377	1.113	0.844	0.566	--
Anti-Hay Mab	0.1 ug/ml	1.016	0.553	0.555	0.402	2.228

*All Group A Streptococcus (GAS) from ATCC (accession numbers noted above); plates were coated with MeOH-fixed bacteria and read at 15 minutes.

EXAMPLE 8

Humanization of the Anti-Staph antibody 96-110 Cloning of the 96-110 variable region cDNAs

The hybridoma cell producing the 96-110 antibody was obtained as described above. A vial of cells was thawed, washed with serum free medium and then resuspended in IMDM (Mediatech) complete media supplemented with 10%FBS (Irvine). Total RNA was isolated from 1×10^8 96-110 cells using the Midi RNA Isolation kit (Qiagen) following the manufacturer's procedure. The RNA was dissolved in 10mM Tris, 0.1mM EDTA (pH8.4) containing 0.03U/ μ g Prime RNase Inhibitor (5'-3') to a final concentration of 0.25 μ g/ μ l.

Figure 10 shows the strategy for cloning the variable region gene fragments and Figure 11 lists the oligonucleotides primers used. The 96-110 total RNA (2 μ g) was converted to cDNA by using Superscript II-MMLV Reverse transcriptase (Life Technologies) and mouse kappa (OKA57) and mouse heavy chain (JS160-162)-specific priming according of manufacturer's procedures. The first strand cDNA synthesis products were then purified using a Centricon-30 concentrator device (Amicon). Of the 40 μ l of cDNA recovered, 5 μ l was used as template DNA for PCR. Typical PCR amplification reactions (100 μ l) contained template DNA, 50 pmoles of the appropriate primers (PMC12-15,55 and OKA57 for light chains, JSS1-4,8 and JS 160-162 for heavy chains), 2.5 units of *ExTaq* polymerase (PanVera), 1x *ExTaq* reaction buffer, 200 μ M dNTP, 1mM $MgCl_2$. The template was denatured by an initial incubation at 96°C for 5 min. The products were amplified by 15 thermal cycles of

cells in an ELISA (see Example 9). Both of these pseudogene transcripts, when converted to cDNA by RT-PCR, contain an Af/III restriction site. For this reason, the PCR products synthesized for the light chain variable region was digested with Af/III and those products that did not cut were then cloned into the pGEM T-Easy (Promega) T/A style cloning vector using the manufacturer's procedures. Light chain candidate (pJRS319) clones were digested with EcoRI (New England Biolabs) using the manufacturer's procedures to identify clones containing inserts of the appropriate size (350bp). The final consensus DNA sequence of the light chain variable regions is shown in Figure 12. The amino acids encoded by these sequences match the N-terminal amino acid analyses of the heavy and light chain peptide fragments produced by the hybridoma cell line.

The heavy and light chain variable regions were then subcloned into a mammalian expression plasmid vector pSUN 15 for production of recombinant chimeric antibody molecules. The creation of the expression vector was an extensive process of DNA fragment ligations and site directed mutagenesis steps. The result was a vector that expresses both antibody chains with CMV promoter driven transcription. Neomycin resistance serves as a dominant selectable marker for transfection of mammalian cells. In addition, it has been designed to allow convenient cloning of any light chain variable region as *EcoRV*/*Bst*BI fragment, any heavy chain variable region as a *Nru*I/*Eco*RI fragment, and any heavy chain constant domain as an *Eco*RI/*Not*I fragment. These restriction sites were chosen because they occur rarely (if ever) in human and mouse variable regions. There is a mouse J region/kappa intron fragment fused to a human

kappa exon so that after post transcriptional splicing a mouse human chimeric kappa light chain is produced.

The backbone of the vector was the plasmid pCDNA3 (Invitrogen). This plasmid was cut with HindIII/XhoI and a "light chain polylinker" DNA fragment was inserted to create the stated "light chain vector." This linker contained the restriction sites HindIII, KpnI, ClaI, PmlI, EcoRV XmaI, BamHI and XhoI to facilitate subsequent cloning steps to create the plasmid pCDNA3.LCPL. A SmaI/ BclI DNA fragment containing a light chain leader, anti-CKMB kappa light chain genomic fragment, and 3' UTR was cloned into the EcoRV/BamHI sites of pCDNA3.LCPL. The mouse kappa intron, exon and 3' UTR in this fragment was derived from LCPXK2 received from Dr. Richard Near (Near, RI et al, 1990, Mol Immunol. 27:901-909). Mutagenesis was then performed to eliminate an NruI (209), MluI (229). and BstBI (2962) and to introduce an NheI (1229) and a BamHI (1214) site to create pCDNA3mut.LCPL.LCVK.

A second "heavy chain vector" was constructed from the pCDNA3mut.LCPL.LCVK plasmid by replacing the light chain expression region (HindIII/XhoI) with a "heavy chain polylinker" consisting of restriction sites HpaI, BspEI, EcoRV, KpnI, and XhoI. A SmaI/ KpnI DNA fragment contains a heavy chain leader, antiCKMB IgG2b chain genomic fragment. A KpnI/SalI oligo nucleotide fragment containing a 3' UTR and a NotI upstream of the SalI site was subsequently cloned into the KpnI1XhoI digested plasmid, (knocking out the XhoI site) to create the plasmid pCDNA3mut.HCPL.HCV2b. From this point two vectors were created that did not have any of the anti-CKMB variable or constant domain DNA sequences. This was done by cutting the plasmid

pCDNA3mut.LCPL.LCVK with EcoRV/XhoI and inserting a linker oligonucleotide fragment containing EcoRV, BstBI, and XhoI sites to create pSUN9. In a similar way, the anti-CKMB fragment in pCDNA3mut.HCPL.HCV2b (NruI/NotI) was replaced by a linker oligonucleotide fragment containing NruI, EcoRI and NotI sites to create pSUN10. A human kappa light chain constant domain was then cloned into pSUN9 as a BstBI/XhoI fragment, and a human IgG1 constant domain was cloned into pSUN10 as a EcoRI/NotI fragment. A BglII/NheI fragment from the human heavy chain vector was then cloned into the human light chain vector cut with BamHI/NheI to create pSUN15. This vector results in the production of recombinant antibody molecules under the control of the CMV transcriptional promoters. The heavy chain molecules are direct cDNA constructs that fuse the variable region sequence directly into the human IgG1 constant domain. The light chain molecules, on the other hand, have a mouse kappa intron region 3' of the variable region coding fragment. After splicing the variable region becomes fused to a human kappa constant region exon. The selectable marker for the vector in mammalian cells is Neomycin (G418).

The variable region gene fragments were re-amplified by PCR using primers that adapted the fragments for cloning into the expression vector (see Figures 12 and 14). The heavy chain front primer (96110HF2) includes a 5' tail that encodes the C-terminus of the heavy chain leader and an *NruI* restriction site for cloning, while the heavy chain reverse primer (96110HB) adds a 3' *EcoRI* restriction site for cloning. The light chain front primer (96110bLF) converts the first amino acid of the 96-110 light chain variable region sequence from glutamine (Q) to aspartic acid (D) via the introduction of an

EcoRV restriction site at the N-terminus of the light chain variable region for cloning, while the light chain reverse primer (96-110bLB) adds a 3' DNA sequence for the joining region-kappa exon splice junction followed by a *BstB1* restriction site for cloning. Because the last amino acid of the light chain variable region is an arginine (R) which is a very rare amino acid at this position, the reverse primer introduces a point mutation in the codon for amino acid 106 that converts it to the much more common lysine (L). This was done because the splice junction in the expression vector for the kappa chain was derived from a J region that encoded a lysine at this position. Neither mutation in the recombinant form of the antibody would be anticipated to alter the antibodies binding characteristics.

PCRs were performed as described above except 10ng of plasmid template was used in each case. Following a 5 min. incubation at 96°C, the PCR parameters were 35 thermal cycles of 58°C for 30 sec., 70°C for 30 sec., and 96°C for 1 min. The 96-110 heavy chain PCR product (approximately 400 bp) was digested with *NruI* and *EcoRI* (New England Biolabs), purified using a Qiaquick PCR Purification column (Qiagen), as described by the manufacturer, and ligated into *NruI* / *EcoRI* digested and gel-purified pSUN15, resulting in plasmid pJRS311 (see Figure 13).

At this point a *BstBI/NotI* (New England Biolabs) DNA fragment containing a mouse kappa J-kappa intron fragment fused to a human kappa exon fragment was digested and gel-purified from the vector tKMC180C1. This fragment was ligated into the backbone of pJRS311 digested with *BstBI/NotI* and gel-purified resulting in the plasmid pJRS315 (see Figure 13).

This was the plasmid into which was cloned the 96-110 light chain variable region. The 96-110 light chain PCR product (approximately 350 bp) was digested with *EcoRV* and *BstBI* (New England Biolabs), purified using a Qiaquick PCR Purification column (Qiagen), as described by the manufacturer, and ligated into *EcoRV/Bs/BI* digested and gel-purified pJRS315, resulting in plasmid pJRS326 (see Figure 13).

It was determined that during this cloning process, a deletion of approximately 200bp occurred at the intron exon junction of the kappa light chain. To repair this, an identical DNA fragment (also a *BstBI/NotI* restriction fragment) was gel-purified from digested pEN22 and ligated into *BstBI/NotI* digested and gel-purified pJRS326, resulting in the final expression plasmid construct pJRS334 (see Figures 13 and 14). The sequence of the variable regions and leader and other junctions was verified prior to mammalian cell transfection.

EXAMPLE 9

Transient production of recombinant chimeric mouse/human 96-110 antibody

Two individual clones of the plasmid pJRS334 (pJRS334-1, -2) were transfected into COS and CHO cells using Superfectant (Qiagen) in 6 well tissue culture cells as described by the manufacturer. After three days the supernatant was assayed for the production of "humanized" antibody and for the capability for the expressed antibody to bind to the heat-killed Staph antigen.

Antibody production assays were preformed in 8-well strips from 96-well microtiter plates (Maxisorp F8; Nunc, Inc.) coated at a 1:500 dilution with Goat anti-Human IgG antibody (Pierce) using a bicarbonate coating buffer, pH 8.5. The plates are covered with pressure sensitive film (Falcon, Becton Dickinson) and incubated overnight at 4°C. Plates are then washed once with Wash solution (Imadazole/NaCl/0.4% Tween-20). 100 microliters of culture supernatant was then applied to duplicate wells and allowed to incubate for 30 minutes on plate rotator at room temperatures. The plates were washed five times with Wash solution. A Goat anti Human kappa-HRP (Southern Biotechnologies) conjugate was diluted 1:800 in the sample/conjugate diluent. 100 microliters was added to the samples, then incubated on a plate rotator for 30 minutes at room temperature. The samples were washed as above and then incubated with 100 µL/well of ABTS developing substrate (Kirkgaard & Perry Laboratories) for 10-15 minutes on a plate rotator at room temperature. The reaction was stopped with 100 µL/well of Quench buffer (Kirkgaard & Perry Laboratories) and the absorbance value at 405 nm was determined using an automated microtiter plate ELISA reader (Ceres UV900HI, Biotek, Winooski, Vermont). As a positive control, a humanized mouse/human chimeric antibody BC24 was used. This assay (see Figure 15) demonstrates that the transfection of cells with this plasmid construct to results in the cells producing a molecule containing both human IgG and kappa domains. The supernatants were then assayed for the ability of the expressed antibodies to bind to heat-killed Staph. The activity assays were preformed in 8-well strips from 96-well microtiter plates (Maxisorp F8; Nunc, Inc.) coated at 0.09 OD/well

with heat-killed Staph Hay cell material suspended in MeOH. The plates are left uncovered and incubated overnight at 4°C. Plates are then washed once PBS. 100 microliters of culture supernatant was then applied to duplicate wells and allowed to incubate for 60 minutes on plate rotator at room temperature. The plates were washed five times with Wash solution. The goat anti Human kappa-HRP was diluted 1:800 in the sample/conjugate diluent. 100 microliters was added to the samples, then incubated on a plate rotator for 30 minutes at room temperatures. The samples were washed as above and then incubated with 100 µL/well of ABTS developing substrate (Kirkgaard & Perry Laboratories) for 10-15 minutes on a plate rotator at room temperature. The reaction was stopped with 100 µL/well of Quench buffer (Kirkgaard & Perry Laboratories) and the absorbance value at 405 nm was determined using an automated microtiter plate ELISA reader (Ceres UV900HI, Bioteck, Winooski, Vermont). As a positive control, the original mouse monoclonal antibody 96-110 was used, and assayed with a Goat anti-Mouse Fc-HRP conjugate @ 1:2000 dilution. This assay (see Figure 16) demonstrates that the transfection of cells with this plasmid construct to results in the cells producing a molecule that binds to the Staph Hay cellular antigen.

EXAMPLE 10

Stable production of recombinant chimeric mouse/human 96-110 antibody

The plasmid pJRS334-1 was transfected into NS/0 cells (obtainable from Baxter International) and CHO cells using electroporation. The plasmid was linearized with

PvuI restriction digestion. 25 micrograms of digested plasmid DNA was mixed with 1×10^7 cells in a total volume of 800 microliters in a 4 centimeter cuvette and subjected to a pulse of 250mA, 9600microF. The cells were plated out after 24 hours in 10ml non-selective media. The cells were then diluted out into 96-well microtiter plates. As colonies appeared, the supernatants were assayed for the production of "humanized" antibody and for the capability for the expressed antibody to bind to the heat-killed Staph antigen. Antibody production and activity assays for the stable transfectants were performed as described above. These assays demonstrate that the transfection of cells with this plasmid construct can result in the production of a stable cell line that produces a humanized chimeric version of the 96-110 mouse hybridoma antibody.

EXAMPLE 11

Opsonic Activity

Having produced a chimeric anti-LTA MAB for staphylococci, we tested its functional activity using *S. epidermidis* as a representative staphylococcal organism. Using the neutrophil mediated opsonophagocytic assay described generally in the Material and Methods section, we assessed the MAB's opsonic activity by evaluating the percent of bacteria killed after two hours of incubation.

Neutrophils, specifically polymorphonuclear neutrophils, were isolated from adult venous blood by dextran sedimentation and ficoll-hypaque density centrifugation. Washed neutrophils were added to round-bottomed wells of microtiter plates (approximately 10^6 cells per well) with approximately 3×10^4 mid-log phase bacteria (*S.*

epidermidis Hay, ATCC 55133). Human sera (10 uls), screened to assure absence of antibody to *S. epidermidis*, was used as a source of active complement (C-Barb-Ex (1:4)).

Forty microliters of immunoglobulin were added at various concentrations (20 ug/ml, 40 ug/ml, 80 ug/ml, and 160 ug/ml) and the plates were incubated at 37°C with constant, vigorous shaking. Samples of 10 uls were taken from each well at zero time and after 2 hours of incubation. Each was diluted, vigorously vortexed to disperse the bacteria, and cultured on blood agar plates overnight at 37°C to quantitate the number of viable bacteria. Results are presented in Figure 17 as percent reduction in numbers of bacterial colonies observed compared to control samples.

Compared to PMN alone or PMN plus complement, the addition of the MAB markedly enhanced opsonic activity for staphylococcus at 20-160 ug/ml). These data demonstrate that the MAB has functional activity and can enhance the phagocytosis and killing of staphylococcal organisms, as represented by *S. epidermidis*.

EXAMPLE 12

In vivo Protective Efficacy

Using the lethal staphylococcal sepsis in adult mice assay (described in Example 3), we compared protection between the original mouse MAB and the chimeric HuMAB. Adult CF1 mice were given 0.5 ml *S. epidermidis* (Hay) IP (3.5×10^9 bacteria). About 24 hrs and 1 hr before infection, 14 mg/kg of each MAB was given to a group of mice,

with a third group of mice given only PBS. All animals were followed for 40 hours after challenge to determine survival.

As set forth in Figure 18, approximately 18 hours after infection, all the control animals died while both treatment groups exhibited 100% survival. At 30 hours after infection, both MAB treatment groups exhibited 70% survival. At the end of the study, the group that received the mouse MAB exhibited greater survival than the group receiving the chimeric MAB, but both MAB enhanced survival over the PBS controls.

We conducted further studies with the chimeric MAB at a dose of 18 mg/kg/dose 2 doses given IP 24 and 1 hour prior to infection (3×10^9 IP *S. epidermidis*, Hay). As set forth in Figure 19, the chimeric MAB enhanced survival.

We also assessed the effect of the chimeric MAB on bacteremia in the lethal *S. epidermidis* sepsis model. CF-1 mice were twice infected IP with strain Hay and the chimeric MAB. Bacteremia is expressed as the number of bacteria isolated on blood agar after a 1:1000 dilutions. As set forth in Figure 20, the chimeric MAB reduced bacterial levels by over 2 logs. Additional studies demonstrated that bacteremia was reduced to a greater degree using 40 mg/kg/dose compared to 20 mg/kg/dose even if survival was comparable. See Figure 21.

These data indicate that increasing the amount of antibody resulted in increased bacterial clearance *in vivo*. Such a response is similar to the observed enhanced opsonic activity *in vitro* as seen when antibody was increased from 20 ug/mg to 160 ug/ml in the neutrophil mediated opsonophagocytic assay (Figure 17).

EXAMPLE 13

In vivo Protective Efficacy

The effect of the chimeric MAB 96-110 was then analyzed in a neonatal staphylococcal model using suckling rats with a foreign body infection. Two day old Wistar rats were treated with lipid emulsion (as is standard in newborn care for nutritional purposes) 0.2 ml, 20% IP on day -1 and again on day +1 and +2 to induce further compromise of the immuno system. In two studies, we injected approximately 5×10^7 of four different strains of *S. epidermidis*, identified below in Table 11 SQ through a plastic catheter and the catheter was left in place under the skin. Saline, 0.2 ml, or MAB 96-110, 0.2 ml (dose of 50-60 mg/kg), was given IP 30 min before and 24 hours after infection. The animals were followed for 5 days.

As set forth in Table 11, in study I, survival for animals receiving MAB ranged from 67% to 83%, with an average of 76%, in contrast to saline treatment, which ranged from 33% to 50%, with an average of 39%. Study II showed even more impressive results. Survival for animals treated with MAB ranged from 83% to 100%, with 90% average, compared to the saline controls at 33% to 50%, with an average of 40%. The complied data for study II are shown in Figure 22.

The Effect of Hu96-110 on Survival in a Lethal Neonatal *S. epidermidis* Sepsis Model

[illegible]

These data demonstrate that the chimeric human antibody directed against LTA is opsonic and enhances survival against staphylococci. In addition, the antibody promotes clearance of the staphylococci from the blood. Thus antibody to LTA provides prophylactic and therapeutic capabilities against staphylococcal infections and vaccines using LTA or peptide mimotopes of LTA that induce anti-LTA antibodies would also have prophylactic capabilities.

Having now fully described the invention, it will be appreciated by those skilled in the art that the invention can be performed within a range of equivalents and conditions without departing from the spirit and scope of the invention and without undue experimentation. In addition, while the invention has been described in light of certain embodiments and examples, the inventors believe that it is capable of further modifications. This application is intended to cover any variations, uses, or adaptations of the invention which follow the general principles set forth above.

The specification includes recitation to the literature and those literature references are herein specifically incorporated by reference.

The specification and examples are exemplary only with the particulars of the claimed invention set forth as follows:

SEQUENCE LISTING

(1) GENERAL INFORMATION:

(i) APPLICANT: Fischer, Gerald W.
Schuman, Richard F.
Wong, Hing
Stinson, Jeffrey L.

(ii) TITLE OF INVENTION: OPSONIC AND PROTECTIVE MONOCLONAL AND
CHIMERIC ANTIBODIES SPECIFIC FOR LIPOTEICHOIC ACID OF GRAM
POSITIVE BACTERIA

(iii) NUMBER OF SEQUENCES: 105

(iv) CORRESPONDENCE ADDRESS:

(A) ADDRESSEE: FINNEGAN, HENDERSON, FARABOW, GARRETT &
DUNNER, LLP
(B) STREET: 1300 I Street, NW
(C) CITY: Washington
(D) STATE: DC
(E) COUNTRY: USA
(F) ZIP: 20005-3315

(v) COMPUTER READABLE FORM:

(A) MEDIUM TYPE: Floppy disk
(B) COMPUTER: IBM PC compatible
(C) OPERATING SYSTEM: PC-DOS/MS-DOS
(D) SOFTWARE: PatentIn Release #1.0, Version #1.30

(vi) CURRENT APPLICATION DATA:

(A) APPLICATION NUMBER: US Unassigned
(B) FILING DATE: Concurrently Herewith

(viii) ATTORNEY/AGENT INFORMATION:

(A) NAME: Einaudi, Carol P.
(B) REGISTRATION NUMBER: 32,220
(C) REFERENCE/DOCKET NUMBER: 04995.0041-00000

(ix) TELECOMMUNICATION INFORMATION:

(A) TELEPHONE: 202-408-4000
(B) TELEFAX: 202-408-4400

04995.0041-00000

(2) INFORMATION FOR SEQ ID NO:1:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 15 amino acids
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS:
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

Trp	Arg	Met	Tyr	Phe	Ser	His	Arg	His	Ala	His	Leu	Arg	Ser	Pro
1				5					10					15

(2) INFORMATION FOR SEQ ID NO:2:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 15 amino acids
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS:
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

Trp	His	Trp	Arg	His	Arg	Ile	Pro	Leu	Gln	Leu	Ala	Ala	Gly	Arg
1				5					10					15

(2) INFORMATION FOR SEQ ID NO:3:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 21 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: other nucleic acid
(A) DESCRIPTION: /desc = "primer"

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:3:

TGAATTTTCT GTATGAGGTT T

21

(2) INFORMATION FOR SEQ ID NO:4:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 30 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(ix) FEATURE:

- (A) NAME/KEY: CDS
- (B) LOCATION: 1..30

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:4:

GGG GCT CAT GCG GAT AGG GTT TAT GGG GCC
Gly Ala His Ala Asp Arg Val Tyr Gly Ala
1 5 10

30

(2) INFORMATION FOR SEQ ID NO:5:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 10 amino acids
 - (B) TYPE: amino acid
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:5:

Gly Ala His Ala Asp Arg Val Tyr Gly Ala
1 5 10

(2) INFORMATION FOR SEQ ID NO:6:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 30 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(ix) FEATURE:

- (A) NAME/KEY: CDS
- (B) LOCATION: 1..30

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:6:

GGG ANT CAT GCG GAT AGG GTT TAT GGG GCC
Gly Xaa His Ala Asp Arg Val Tyr Gly Ala
1 5 10

30

(2) INFORMATION FOR SEQ ID NO:7:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 10 amino acids
- (B) TYPE: amino acid
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:7:

Gly Xaa His Ala Asp Arg Val Tyr Gly Ala
1 5 10

(2) INFORMATION FOR SEQ ID NO:8:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 57 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

- (ix) FEATURE:
 (A) NAME/KEY: CDS
 (B) LOCATION: 1..57

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:8:

GGG GCT TGG CAT TGG CGT CAT CGT ATT CCT CTT CAG CTT GCT GCT GGT 48
Gly Ala Trp His Trp Arg His Arg Ile Pro Leu Gln Leu Ala Ala Gly
1 5 10 15

CGT GGG GCC 57
Arg Gly Ala

(2) INFORMATION FOR SEQ ID NO:9:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 19 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:9:

Gly Ala Trp His Trp Arg His Arg Ile Pro Leu Gln Leu Ala Ala Gly
1 5 10 15

Arg Gly Ala

(2) INFORMATION FOR SEQ ID NO:10:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 57 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

- (ix) FEATURE:
 (A) NAME/KEY: CDS
 (B) LOCATION: 1..57

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:10:

GGG GCT CGT CGG CAT GGT AAT TTT TCT CAT TTT TTT CAT CGG TCG TTG 48
Gly Ala Arg Arg His Gly Asn Phe Ser His Phe Phe His Arg Ser Leu
 1 5 10 15

ATT GGG GCC 57
Ile Gly Ala

(2) INFORMATION FOR SEQ ID NO:11:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 19 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:11:

Gly Ala Arg Arg His Gly Asn Phe Ser His Phe Phe His Arg Ser Leu
 1 5 10 15

Ile Gly Ala

(2) INFORMATION FOR SEQ ID NO:12:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 57 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

- (ix) FEATURE:
 (A) NAME/KEY: CDS
 (B) LOCATION: 1..57

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:12:

GGG GCT TGG AAG GCT TTG TTT AGT CAT TCT TAT CGT CCT CGG GGT TCG 48
Gly Ala Trp Lys Ala Leu Phe Ser His Ser Tyr Arg Pro Arg Gly Ser
 1 5 10 15

GCT GGG GCC 57
Ala Gly Ala

(2) INFORMATION FOR SEQ ID NO:13:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 19 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:13:

Gly Ala Trp Lys Ala Leu Phe Ser His Ser Tyr Arg Pro Arg Gly Ser
 1 5 10 15

Ala Gly Ala

(2) INFORMATION FOR SEQ ID NO:14:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 57 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(ix) FEATURE:

- (A) NAME/KEY: CDS
- (B) LOCATION: 1..57

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:14:

GGG GCT AGG CAT TGG CGT CAT CGT ATT CCT CTT CAG CTT GCT GCT GGT 48
Gly Ala Arg His Trp Arg His Arg Ile Pro Leu Gln Leu Ala Ala Gly
1 5 10 15
CGT GGG GCC 57
Arg Gly Ala

(2) INFORMATION FOR SEQ ID NO:15:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 19 amino acids
- (B) TYPE: amino acid
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:15:

Gly Ala Arg His Trp Arg His Arg Ile Pro Leu Gln Leu Ala Ala Gly
1 5 10 15
Arg Gly Ala

(2) INFORMATION FOR SEQ ID NO:16:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 57 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

- (ix) FEATURE:
 (A) NAME/KEY: CDS
 (B) LOCATION: 1..57

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:16:

GGG GCT TGG CAT TGG CGT CAT CGT ATT CCT CTT CAG CTT GCT GCT GGT 48
Gly Ala Trp His Trp Arg His Arg Ile Pro Leu Gln Leu Ala Ala Gly
 1 5 10 15

CGT GGG GCC 57
Arg Gly Ala

(2) INFORMATION FOR SEQ ID NO:17:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 19 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:17:

Gly Ala Trp His Trp Arg His Arg Ile Pro Leu Gln Leu Ala Ala Gly
 1 5 10 15

Arg Gly Ala

(2) INFORMATION FOR SEQ ID NO:18:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 57 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

- (ix) FEATURE:
 (A) NAME/KEY: CDS
 (B) LOCATION: 1..57

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:18:

GGG GCT TGG CAT TGG CGT CAT CGT ATT CCT CTT CAG CTT GCT GCT GGT 48
Gly Ala Trp His Trp Arg His Arg Ile Pro Leu Gln Leu Ala Ala Gly
 1 5 10 15

CGT GGG GCC 57
Arg Gly Ala

(2) INFORMATION FOR SEQ ID NO:19:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 19 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:19:

Gly Ala Trp His Trp Arg His Arg Ile Pro Leu Gln Leu Ala Ala Gly
 1 5 10 15

Arg Gly Ala

(2) INFORMATION FOR SEQ ID NO:20:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 57 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

- (ix) FEATURE:
 (A) NAME/KEY: CDS
 (B) LOCATION: 1..57

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:20:

GGG GCT CAG GTG GCT GTT TTG TAT CCT CCT TTG GCT GAT GCT ACT GAG 48
Gly Ala Gln Val Ala Val Leu Tyr Pro Pro Leu Ala Asp Ala Thr Glu
 1 5 10 15

CTT GGG GCC 57
Leu Gly Ala

(2) INFORMATION FOR SEQ ID NO:21:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 19 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:21:

Gly Ala Gln Val Ala Val Leu Tyr Pro Pro Leu Ala Asp Ala Thr Glu
 1 5 10 15

Leu Gly Ala

Table 1. *Continued*

Parameter	Unit	Value	Standard error	95% CI
Intercept		1.00	0.00	1.00
Age	yr	0.01	0.01	-0.01, 0.03
Gender				
Male		0.00	0.01	-0.02, 0.02
Female		0.00	0.01	-0.02, 0.02
Marital status				
Married		0.00	0.01	-0.02, 0.02
Single		0.00	0.01	-0.02, 0.02
Divorced		0.00	0.01	-0.02, 0.02
Widowed		0.00	0.01	-0.02, 0.02
Education	yr	0.00	0.01	-0.02, 0.02
Income	\$1000	0.00	0.01	-0.02, 0.02
Health status				
Excellent		0.00	0.01	-0.02, 0.02
Very good		0.00	0.01	-0.02, 0.02
Good		0.00	0.01	-0.02, 0.02
Fair		0.00	0.01	-0.02, 0.02
Poor		0.00	0.01	-0.02, 0.02
Chronic conditions				
Hypertension		0.00	0.01	-0.02, 0.02
Diabetes		0.00	0.01	-0.02, 0.02
Asthma		0.00	0.01	-0.02, 0.02
Heart disease		0.00	0.01	-0.02, 0.02
Stroke		0.00	0.01	-0.02, 0.02
Cancer		0.00	0.01	-0.02, 0.02
Other		0.00	0.01	-0.02, 0.02
Medication				
None		0.00	0.01	-0.02, 0.02
One or more		0.00	0.01	-0.02, 0.02
Physical activity				
Sedentary		0.00	0.01	-0.02, 0.02
Light		0.00	0.01	-0.02, 0.02
Moderate		0.00	0.01	-0.02, 0.02
Vigorous		0.00	0.01	-0.02, 0.02
Stress				
Low		0.00	0.01	-0.02, 0.02
Medium		0.00	0.01	-0.02, 0.02
High		0.00	0.01	-0.02, 0.02
Life satisfaction				
Very satisfied		0.00	0.01	-0.02, 0.02
Satisfied		0.00	0.01	-0.02, 0.02
Dissatisfied		0.00	0.01	-0.02, 0.02
Very dissatisfied		0.00	0.01	-0.02, 0.02

(ii) MOLECULE TYPE: DNA

- (ix) FEATURE:

- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:22:

GGG GCT CGT CGG CAT GGT AAT TTT TCT CAT TTT TTT CAT CGG TCG TTG	48
Gly Ala Arg Arg His Gly Asn Phe Ser His Phe Phe His Arg Ser Leu	
1 5 10 15	
ATT GGG GCC	57
Ile Gly Ala	

(i) SEQUENCE CHARACTERISTICS:

- (ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:23:

Gly Ala Arg Arg His Gly Asn Phe Ser His Phe Phe His Arg Ser Leu
1 5 10 15
Ile Gly Ala

(2) INFORMATION FOR SEQ ID NO:24:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 57 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

- (ix) FEATURE:
 (A) NAME/KEY: CDS
 (B) LOCATION: 1..57

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:24:

GGG GCT CGT CGG CAT GGT AAT TTT TCT CAT TTT TTT CAT CGG TCG TTG 48
Gly Ala Arg Arg His Gly Asn Phe Ser His Phe Phe His Arg Ser Leu
 1 5 10 15

ATT GGG GCC 57
Ile Gly Ala

(2) INFORMATION FOR SEQ ID NO:25:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 19 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:25:

Gly Ala Arg Arg His Gly Asn Phe Ser His Phe Phe His Arg Ser Leu
 1 5 10 15

Ile Gly Ala

(2) INFORMATION FOR SEQ ID NO:26:

- (i) SEQUENCE CHARACTERISTICS:

	Mean	SD	Median	Mode	Range	Skewness	Kurtosis
Age (years)	45.2	10.5	42.0	40.0	25-65	0.15	2.10
Gender (male/female)	50/50						
Education (years)	12.5	2.5	12.0	12.0	8-16	0.20	2.20
Occupation (white/blue)	55/45						
Marital status (married/divorced)	60/40						
Income (thousands \$)	25.0	15.0	20.0	20.0	10-50	0.30	2.30
Health status (good/fair/poor)	40/30/30						
Smoking status (smoker/non-smoker)	30/70						
Alcohol consumption (drinks/week)	2.0	1.5	1.0	1.0	0-5	0.10	2.00
Exercise frequency (times/week)	1.0	1.0	0.0	0.0	0-3	0.20	2.10
Stress level (low/moderate/high)	30/40/30						
Family size (children)	2.0	1.0	1.0	1.0	0-4	0.10	2.00
Home ownership (owner/renter)	65/35						
Commute time (minutes)	20.0	10.0	15.0	15.0	5-45	0.20	2.10
Neighborhood safety (safe/unsafe)	70/30						
Access to healthcare (yes/no)	90/10						
Health insurance (has/doesn't)	95/5						
Chronic conditions (0-5)	1.5	1.0	1.0	1.0	0-5	0.10	2.00
Medication use (yes/no)	40/60						
Healthcare costs (thousands \$)	10.0	5.0	8.0	8.0	0-20	0.20	2.10
Healthcare satisfaction (satisfied/dissatisfied)	50/50						
Healthcare access (easy/difficult)	40/60						
Healthcare quality (good/bad)	55/45						
Healthcare wait time (minutes)	30.0	15.0	25.0	25.0	10-60	0.20	2.10
Healthcare cost (thousands \$)	10.0	5.0	8.0	8.0	0-20	0.20	2.10
Healthcare quality (good/bad)	55/45						
Healthcare wait time (minutes)	30.0	15.0	25.0	25.0	10-60	0.20	2.10
Healthcare cost (thousands \$)	10.0	5.0	8.0	8.0	0-20	0.20	2.10
Healthcare quality (good/bad)	55/45						
Healthcare wait time (minutes)	30.0	15.0	25.0	25.0	10-60	0.20	2.10
Healthcare cost (thousands \$)	10.0	5.0	8.0	8.0	0-20	0.20	2.10
Healthcare quality (good/bad)	55/45						
Healthcare wait time (minutes)	30.0	15.0	25.0	25.0	10-60	0.20	2.10
Healthcare cost (thousands \$)	10.0	5.0	8.0	8.0	0-20	0.20	2.10
Healthcare quality (good/bad)	55/45						
Healthcare wait time (minutes)	30.0	15.0	25.0	25.0	10-60	0.20	2.10
Healthcare cost (thousands \$)	10.0	5.0	8.0	8.0	0-20	0.20	2.10
Healthcare quality (good/bad)	55/45						
Healthcare wait time (minutes)	30.0	15.0	25.0	25.0	10-60	0.20	2.10
Healthcare cost (thousands \$)	10.0	5.0	8.0	8.0	0-20	0.20	2.10
Healthcare quality (good/bad)	55/45						
Healthcare wait time (minutes)	30.0	15.0	25.0	25.0	10-60	0.20	2.10
Healthcare cost (thousands \$)	10.0	5.0	8.0	8.0	0-20	0.20	2.10
Healthcare quality (good/bad)	55/45						
Healthcare wait time (minutes)	30.0	15.0	25.0	25.0	10-60	0.20	2.10
Healthcare cost (thousands \$)	10.0	5.0	8.0	8.0	0-20	0.20	2.10
Healthcare quality (good/bad)	55/45						
Healthcare wait time (minutes)	30.0	15.0	25.0	25.0	10-60	0.20	2.10
Healthcare cost (thousands \$)	10.0	5.0	8.0	8.0	0-20	0.20	2.10
Healthcare quality (good/bad)	55/45						
Healthcare wait time (minutes)	30.0	15.0	25.0	25.0	10-60	0.20	2.10
Healthcare cost (thousands \$)	10.0	5.0	8.0	8.0	0-20	0.20	2.10
Healthcare quality (good/bad)	55/45						

(ix) FEATURE:

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:26:

CCT GGG GCC
Pro Gly Ala

57

(i) SEQUENCE CHARACTERISTICS:

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:27:

Pro Gly Ala

(2) INFORMATION FOR SEQ ID NO:28:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 57 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(ix) FEATURE:

- (A) NAME/KEY: CDS
- (B) LOCATION: 1..57

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:28:

GGG GCT TGG CGT ATG TAT TTT TCT CAT CGT CAT GCG CAT CTT CGT AGT	48
Gly Ala Trp Arg Met Tyr Phe Ser His Arg His Ala His Leu Arg Ser	
1 5 10 15	
CCT GGG GCC	
Pro Gly Ala	57

(2) INFORMATION FOR SEQ ID NO:29:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 19 amino acids
- (B) TYPE: amino acid
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:29:

Gly Ala Trp Arg Met Tyr Phe Ser His Arg His Ala His Leu Arg Ser	
1 5 10 15	
Pro Gly Ala	

(2) INFORMATION FOR SEQ ID NO:30:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 57 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(ix) FEATURE:

- (A) NAME/KEY: CDS
- (B) LOCATION: 1..57

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:30:

GGG GCT TGG CGG AAG TAT TTT TCT TAT CAT CAT GCG CAT CTT TGT AGT	48
Gly Ala Trp Arg Lys Tyr Phe Ser Tyr His His Ala His Leu Cys Ser	
1 5 10 15	
CCT GGG GCC	57
Pro Gly Ala	

(2) INFORMATION FOR SEQ ID NO:31:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 19 amino acids
- (B) TYPE: amino acid
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:31:

Gly Ala Trp Arg Lys Tyr Phe Ser Tyr His His Ala His Leu Cys Ser	
1 5 10 15	
Pro Gly Ala	

(2) INFORMATION FOR SEQ ID NO:32:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 57 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

- (ix) FEATURE:
 (A) NAME/KEY: CDS
 (B) LOCATION: 1..57

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:32:

GGG GCT TGG CGT ATG TAT TTT TCT CAT CGT CAT GCG CAT CTT CGT AGT 48
Gly Ala Trp Arg Met Tyr Phe Ser His Arg His Ala His Leu Arg Ser
 1 5 10 15

CCT GGG GCC 57
Pro Gly Ala

(2) INFORMATION FOR SEQ ID NO:33:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 19 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:33:

Gly Ala Trp Arg Met Tyr Phe Ser His Arg His Ala His Leu Arg Ser
 1 5 10 15

Pro Gly Ala

(2) INFORMATION FOR SEQ ID NO:34:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 57 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(ix) FEATURE:

- (A) NAME/KEY: CDS
- (B) LOCATION: 1..57

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:34:

GGG GCT TGG CGT ATG TAT TTT TCT CAT CGT CAT GCG CAT CTT CGT AGT 48
Gly Ala Trp Arg Met Tyr Phe Ser His Arg His Ala His Leu Arg Ser
1 5 10 15
CCT GGG GCC
Pro Gly Ala 57

(2) INFORMATION FOR SEQ ID NO:35:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 19 amino acids
- (B) TYPE: amino acid
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:35:

Gly Ala Trp Arg Met Tyr Phe Ser His Arg His Ala His Leu Arg Ser
1 5 10 15
Pro Gly Ala

(2) INFORMATION FOR SEQ ID NO:36:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 57 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

- (ix) FEATURE:
 (A) NAME/KEY: CDS
 (B) LOCATION: 1..57

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:36:

GGG GCT TGG CGT ATG TAT TTT TCT CAT CGT CAT GCG CAT CTT CGT AGT	48
Gly Ala Trp Arg Met Tyr Phe Ser His Arg His Ala His Leu Arg Ser	
1 5 10 15	
CCT GGG GCC	57
Pro Gly Ala	

(2) INFORMATION FOR SEQ ID NO:37:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 19 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:37:

Gly Ala Trp Arg Met Tyr Phe Ser His Arg His Ala His Leu Arg Ser	
1 5 10 15	
Pro Gly Ala	

[illegible]

(A) LENGTH: 57 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(A) NAME/KEY: CDS
(B) LOCATION: 1..57

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:38:

[illegible]

(2) INFORMATION FOR SEQ ID NO:39:

(A) LENGTH: 19 amino acids
(B) TYPE: amino acid
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:39:

Gly Ala Arg Arg His Gly Asn Phe Ser His Phe Phe His Arg Ser Leu
1 5 10 15
Ile Gly Ala

	1970	1971	1972	1973	1974	1975	1976	1977	1978	1979	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	2022	2023	2024	2025	2026	2027	2028	2029	2030	2031	2032	2033	2034	2035	2036	2037	2038	2039	2040	2041	2042	2043	2044	2045	2046	2047	2048	2049	2050	2051	2052	2053	2054	2055	2056	2057	2058	2059	2060	2061	2062	2063	2064	2065	2066	2067	2068	2069	2070	2071	2072	2073	2074	2075	2076	2077	2078	2079	2080	2081	2082	2083	2084	2085	2086	2087	2088	2089	2090	2091	2092	2093	2094	2095	2096	2097	2098	2099	2100	2101	2102	2103	2104	2105	2106	2107	2108	2109	2110	2111	2112	2113	2114	2115	2116	2117	2118	2119	2120	2121	2122	2123	2124	2125	2126	2127	2128	2129	2130	2131	2132	2133	2134	2135	2136	2137	2138	2139	2140	2141	2142	2143	2144	2145	2146	2147	2148	2149	2150	2151	2152	2153	2154	2155	2156	2157	2158	2159	2160	2161	2162	2163	2164	2165	2166	2167	2168	2169	2170	2171	2172	2173	2174	2175	2176	2177	2178	2179	2180	2181	2182	2183	2184	2185	2186	2187	2188	2189	2190	2191	2192	2193	2194	2195	2196	2197	2198	2199	2200	2201	2202	2203	2204	2205	2206	2207	2208	2209	2210	2211	2212	2213	2214	2215	2216	2217	2218	2219	2220	2221	2222	2223	2224	2225	2226	2227	2228	2229	2230	2231	2232	2233	2234	2235	2236	2237	2238	2239	2240	2241	2242	2243	2244	2245	2246	2247	2248	2249	2250	2251	2252	2253	2254	2255	2256	2257	2258	2259	2260	2261	2262	2263	2264	2265	2266	2267	2268	2269	2270	2271	2272	2273	2274	2275	2276	2277	2278	2279	2280	2281	2282	2283	2284	2285	2286	2287	2288	2289	2290	2291	2292	2293	2294	2295	2296	2297	2298	2299	2300	2301	2302	2303	2304	2305	2306	2307	2308	2309	2310	2311	2312	2313	2314	2315	2316	2317	2318	2319	2320	2321	2322	2323	2324	2325	2326	2327	2328	2329	2330	2331	2332	2333	2334	2335	2336	2337	2338	2339	2340	2341	2342	2343	2344	2345	2346	2347	2348	2349	2350	2351	2352	2353	2354	2355	2356	2357	2358	2359	2360	2361	2362	2363	2364	2365	2366	2367	2368	2369	2370	2371	2372	2373	2374	2375	2376	2377	2378	2379	2380	2381	2382	2383	2384	2385	2386	2387	2388	2389	2390	2391	2392	2393	2394	2395	2396	2397	2398	2399	2400	2401	2402	2403	2404	2405	2406	2407	2408	2409	2410	2411	2412	2413	2414	2415	2416	2417	2418	2419	2420	2421	2422	2
--	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	---

(A) LENGTH: 57 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ix) FEATURE:

- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:40:

[illegible]

(i) SEQUENCE CHARACTERISTICS:

- (ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:41:

Gly Ala Trp His Trp Arg His Arg Ile Pro Leu Gln Leu Ala Ala Gly
1 5 10 15
Arg Gly Ala

(2) INFORMATION FOR SEQ ID NO:42:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 57 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

- (ix) FEATURE:
 (A) NAME/KEY: CDS
 (B) LOCATION: 1..57

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:42:

GGG GCT CGT CGG CAT GGT AAT TTT TCT CAT TTT TTT CAT CGG TCG TTG 48
Gly Ala Arg Arg His Gly Asn Phe Ser His Phe Phe His Arg Ser Leu
 1 5 10 15

ATT GGG GCC 57
Ile Gly Ala

(2) INFORMATION FOR SEQ ID NO:43:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 19 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:43:

Gly Ala Arg Arg His Gly Asn Phe Ser His Phe Phe His Arg Ser Leu
 1 5 10 15

Ile Gly Ala

[illegible]

(ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:44:

[illegible]

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 19 amino acids
- (B) TYPE: amino acid
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:45:

Gly Ala Asp Trp Ile Thr Phe His Arg Arg His His Asp Arg Val Leu
1 5 10 15
Ser Gly Ala

(2) INFORMATION FOR SEQ ID NO:46:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 57 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

- (ix) FEATURE:
 (A) NAME/KEY: CDS
 (B) LOCATION: 1..57

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:46:

GGG GCT GGT TGG ATT ACT TTT CAT CGT CGT CAT CAT GAT CGT GTT CTT	48
Gly Ala Gly Trp Ile Thr Phe His Arg Arg His His Asp Arg Val Leu	
1 5 10 15	
TCT GGG GCC	57
Ser Gly Ala	

(2) INFORMATION FOR SEQ ID NO:47:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 19 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:47:

Gly Ala Gly Trp Ile Thr Phe His Arg Arg His His Asp Arg Val Leu	
1 5 10 15	
Ser Gly Ala	

[illegible]

(A) LENGTH: 57 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ix) FEATURE:

- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:48:

GGG GCT GGG AAG GCT ATG TTT AGT CAT TCT TAT CGT CAT CGG GGT TCG	48
Gly Ala Gly Lys Ala Met Phe Ser His Ser Tyr Arg His Arg Gly Ser	
1 5 10 15	
 GCT GGG GCC	57
Ala Gly Ala	

(i) SEQUENCE CHARACTERISTICS:

- (ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:49:

Gly Ala Gly Lys Ala Met Phe Ser His Ser Tyr Arg His Arg Gly Ser
1 5 10 15
Ala Gly Ala

(2) INFORMATION FOR SEQ ID NO:50:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 57 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

- (ix) FEATURE:
 (A) NAME/KEY: CDS
 (B) LOCATION: 1..57

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:50:

GGG GCT GAT TGG ATT ACT TTT CAT CGT CGT CAT CAT GAT CGT GTT CTT 48
Gly Ala Asp Trp Ile Thr Phe His Arg Arg His His Asp Arg Val Leu
 1 5 10 15

TCT GGG GCC 57
Ser Gly Ala

(2) INFORMATION FOR SEQ ID NO:51:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 19 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:51:

Gly Ala Asp Trp Ile Thr Phe His Arg Arg His His Asp Arg Val Leu
 1 5 10 15

Ser Gly Ala

(2) INFORMATION FOR SEQ ID NO:52:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 57 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

- (ix) FEATURE:
 (A) NAME/KEY: CDS
 (B) LOCATION: 1..57

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:52:

GGG GCT AGT CGT CAT ATG CTT GCT CGG TGG TCG CGT TTG CTT GCT GTT 48
Gly Ala Ser Arg His Met Leu Ala Arg Trp Ser Arg Leu Leu Ala Val
 1 5 10 15

CCT GGG GCC 57
Pro Gly Ala

(2) INFORMATION FOR SEQ ID NO:53:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 19 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:53:

Gly Ala Ser Arg His Met Leu Ala Arg Trp Ser Arg Leu Leu Ala Val
 1 5 10 15

Pro Gly Ala

(2) INFORMATION FOR SEQ ID NO:54:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 57 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

- (ix) FEATURE:
 (A) NAME/KEY: CDS
 (B) LOCATION: 1..57

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:54:

GGG GCT GGG AAG GCT ATG TTT AGT CAT TCT TAT CGT CAT CGG GGT TCG 48
Gly Ala Gly Lys Ala Met Phe Ser His Ser Tyr Arg His Arg Gly Ser
 1 5 10 15

GCT GGG GCC 57
Ala Gly Ala

(2) INFORMATION FOR SEQ ID NO:55:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 19 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:55:

Gly Ala Gly Lys Ala Met Phe Ser His Ser Tyr Arg His Arg Gly Ser
 1 5 10 15

Ala Gly Ala

[illegible]

(A) LENGTH: 57 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(A) NAME/KEY: CDS
(B) LOCATION: 1..57

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:56:

[illegible]

(2) INFORMATION FOR SEQ ID NO:57:

(A) LENGTH: 19 amino acids
(B) TYPE: amino acid
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:57:

Gly Ala Trp His Trp Arg His Arg Ile Pro Leu Gln Leu Ala Ala Gly
1 5 10 15

Arg Gly Ala

(2) INFORMATION FOR SEQ ID NO:58:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 57 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

- (ix) FEATURE:
 (A) NAME/KEY: CDS
 (B) LOCATION: 1..57

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:58:

GGG GCT CGT CGG CAT GGT AAT TTT TCT CAT TTT TTT CAT CGG TCG TTG 48
Gly Ala Arg Arg His Gly Asn Phe Ser His Phe Phe His Arg Ser Leu
 1 5 10 15

ATT GGG GCC 57
Ile Gly Ala

(2) INFORMATION FOR SEQ ID NO:59:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 19 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:59:

Gly Ala Arg Arg His Gly Asn Phe Ser His Phe Phe His Arg Ser Leu
 1 5 10 15

Ile Gly Ala

(2) INFORMATION FOR SEQ ID NO:60:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 57 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(ix) FEATURE:

- (A) NAME/KEY: CDS
- (B) LOCATION: 1..57

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:60:

GGG GCT TGG AAG GCT TTG TTT AGT CAT TCT TAT CGT CCT CGG GGT TCG 48
Gly Ala Trp Lys Ala Leu Phe Ser His Ser Tyr Arg Pro Arg Gly Ser
1 5 10 15
GCT GGG GCC 57
Ala Gly Ala

(2) INFORMATION FOR SEQ ID NO:61:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 19 amino acids
- (B) TYPE: amino acid
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:61:

Gly Ala Trp Lys Ala Leu Phe Ser His Ser Tyr Arg Pro Arg Gly Ser
1 5 10 15
Ala Gly Ala

Table 1. Demographic characteristics of the study population	
Age (years)	65.2 ± 1.2
Gender (male/female)	10/10
Education (years)	12.5 ± 0.5
Occupation (white/blue)	10/10
Marital status (married/divorced/widowed)	10/10/0
Smoking status (smoker/nonsmoker)	10/10
Alcohol consumption (yes/no)	10/10
Comorbidities (hypertension/diabetes/cholesterol)	10/10/10
Medication (antihypertensive/antidiabetic/anticholesterol)	10/10/10
Family history (hypertension/diabetes/cholesterol)	10/10/10
Physical activity (yes/no)	10/10
Stress level (low/moderate/high)	10/10/10
Social support (yes/no)	10/10
Quality of life (SF-36)	50.0 ± 5.0
Health status (good/fair/poor)	10/10/10
Life expectancy (years)	15.0 ± 2.0
Healthcare utilization (yes/no)	10/10
Health insurance (yes/no)	10/10
Health literacy (yes/no)	10/10
Health beliefs (yes/no)	10/10
Health behaviors (yes/no)	10/10
Health outcomes (yes/no)	10/10
Health status (good/fair/poor)	10/10/10
Life expectancy (years)	15.0 ± 2.0
Healthcare utilization (yes/no)	10/10
Health insurance (yes/no)	10/10
Health literacy (yes/no)	10/10
Health beliefs (yes/no)	10/10
Health behaviors (yes/no)	10/10
Health outcomes (yes/no)	10/10

```
(A) LENGTH: 57 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear
```

(ix) FEATURE:

- (A) NAME/KEY: CDS
(B) LOCATION: 1..57

GGG GCT CAG GTG GCT GTT TTG TAT CCT CCT TTG GCT GAT GCT ACT GAG 48
Gly Ala Gln Val Ala Val Leu Tyr Pro Pro Leu Ala Asp Ala Thr Glu
1 5 10 15

CTT GGG GCC
Leu Gly Ala

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 19 amino acids
(B) TYPE: amino acid
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:63:

Gly Ala Gln Val Ala Val Leu Tyr Pro Pro Leu Ala Asp Ala Thr Glu
1 5 10 15

Leu Gly Ala

[illegible]

```
(A) LENGTH: 57 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear
```

(ix) FEATURE:

- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:64:

[illegible]

(i) SEQUENCE CHARACTERISTICS:

- (ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:65:

Gly Ala Trp Arg Met Tyr Phe Ser His Arg His Ala His Leu Arg Ser
1 5 10 15
Pro Gly Ala

(2) INFORMATION FOR SEQ ID NO:66:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 30 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

- (ix) FEATURE:
 (A) NAME/KEY: CDS
 (B) LOCATION: 1..30

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:66:

GGG GCT CAT GCG GAT AGG GTT TAT GGG GCC
Gly Ala His Ala Asp Arg Val Tyr Gly Ala
1 5 10

30

(2) INFORMATION FOR SEQ ID NO:67:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 10 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:67:

Gly Ala His Ala Asp Arg Val Tyr Gly Ala
1 5 10

(2) INFORMATION FOR SEQ ID NO:68:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 45 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: other nucleic acid
 (A) DESCRIPTION: /desc = "primer"

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:68:

ATTTCAGGCC CAGCCGGCCA TGGCCGARGT RMAGCTKSAK GAGWC

45

(2) INFORMATION FOR SEQ ID NO:69:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 45 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: other nucleic acid

(A) DESCRIPTION: /desc = "primer"

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:69:

ATTTCAGGCC CAGCCGGCCA TGGCCGARGT YCARCTKCAR CARYC

45

(2) INFORMATION FOR SEQ ID NO:70:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 45 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: other nucleic acid

(A) DESCRIPTION: /desc = "primer"

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:70:

ATTTCAGGCC CAGCCGGCCA TGGCCCAGGT GAAGCTKSTS GARTC

45

(2) INFORMATION FOR SEQ ID NO:71:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 45 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

- (ii) MOLECULE TYPE: other nucleic acid
 - (A) DESCRIPTION: /desc = "primer"

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:71:

ATTTCAGGCC CAGCCGGCCA TGGCCGAVGT GMWGCTKGTG GAGWC

45

(2) INFORMATION FOR SEQ ID NO:72:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 45 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

- (ii) MOLECULE TYPE: other nucleic acid
 - (A) DESCRIPTION: /desc = "primer"

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:72:

ATTTCAGGCC CAGCCGGCCA TGGCCCAGGT BCARCTKMAR SARTC

45

(2) INFORMATION FOR SEQ ID NO:73:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 35 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

- (ii) MOLECULE TYPE: other nucleic acid
 - (A) DESCRIPTION: /desc = "primer"

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:73:

GCTGCCACCG CCACCTGMRG AGACDGTGAS TGARG

35

(2) INFORMATION FOR SEQ ID NO:74:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 35 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: other nucleic acid

- (A) DESCRIPTION: /desc = "primer"

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:74:

GCTGCCACCG CCACCTGMRG AGACDGTGAS MGTRG

35

(2) INFORMATION FOR SEQ ID NO:75:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 35 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: other nucleic acid

- (A) DESCRIPTION: /desc = "primer"

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:75:

GCTGCCACCG CCACCTGMRG AGACDGTGAS CAGRG

35

(2) INFORMATION FOR SEQ ID NO:76:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 32 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: other nucleic acid

- (A) DESCRIPTION: /desc = "primer"

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:76:

CCCGGGCCAC CATGGAGACA GACACACTCC TG

32

(2) INFORMATION FOR SEQ ID NO:77:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 35 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: other nucleic acid

- (A) DESCRIPTION: /desc = "primer"

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:77:

CCCGGGCCAC CATGGATTTT CAAGTGCAGA TTTC

35

(2) INFORMATION FOR SEQ ID NO:78:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 32 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: other nucleic acid

- (A) DESCRIPTION: /desc = "primer"

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:78:

CCCGGGCCAC CATGGAGWCA CAKWCTCAGG TC

32

(2) INFORMATION FOR SEQ ID NO:79:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 33 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: other nucleic acid

(A) DESCRIPTION: /desc = "primer"

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:79:

CCCGGGCCAC CATGKCCCCW RCTCAGYTTC TKG

33

(2) INFORMATION FOR SEQ ID NO:80:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 31 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: other nucleic acid

(A) DESCRIPTION: /desc = "primer"

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:80:

CCCGGGCACC ATGAAGTTGC CTGTTAGGCT G

31

(2) INFORMATION FOR SEQ ID NO:81:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 23 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: other nucleic acid

(A) DESCRIPTION: /desc = "primer"

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:81:

GCACCTCCAG ATGTAACTG CTC

23

(2) INFORMATION FOR SEQ ID NO:82:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 54 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

- (ii) MOLECULE TYPE: other nucleic acid
(A) DESCRIPTION: /desc = "primer"

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:82:

TAATATCGCG ACAGCTACAG GTGTCCACTC CCGAAGTGAT GCTGGTGGAG WCTG

54

(2) INFORMATION FOR SEQ ID NO:83:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 30 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

- (ii) MOLECULE TYPE: other nucleic acid
(A) DESCRIPTION: /desc = "primer"

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:83:

TTATAGAATT CTGAGGAGAC GGTGAGTGAG

30

(2) INFORMATION FOR SEQ ID NO:84:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 29 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

- (ii) MOLECULE TYPE: other nucleic acid
(A) DESCRIPTION: /desc = "primer"

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:84:

TTAGGCGATA TCGTTCTCTC CCA GTCTCC

29

(2) INFORMATION FOR SEQ ID NO:85:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 46 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: other nucleic acid
(A) DESCRIPTION: /desc = "primer"

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:85:

GTAACCGTTC GAAAAGTGTA CTTACGTTTT ATTTCCAGCA TGGTCC

46

(2) INFORMATION FOR SEQ ID NO:86:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 90 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(ix) FEATURE:
(A) NAME/KEY: CDS
(B) LOCATION: 1..90

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:86:

GAA GTG ATG CTG GTG GAG TCT GGT GGA GGA TTG GTG CAG CCT AAA GGG 48
Glu Val Met Leu Val Glu Ser Gly Gly Gly Leu Val Gln Pro Lys Gly
1 5 10 15

TCA TTG AAA CTC TCA TGT GCA GCC TCT GGA TTC ACC TTC AAT 90
Ser Leu Lys Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Asn
20 25 30

(2) INFORMATION FOR SEQ ID NO:87:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 30 amino acids
(B) TYPE: amino acid
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:87:

Glu Val Met Leu Val Glu Ser Gly Gly Gly Leu Val Gln Pro Lys Gly
 1 5 10 15

Ser Leu Lys Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Asn
 20 25 30

(2) INFORMATION FOR SEQ ID NO:88:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 57 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(ix) FEATURE:

- (A) NAME/KEY: CDS
- (B) LOCATION: 1..57

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:88:

AAC TAC GCC ATG AAT TGG GTC CGC CAG GCT CCA GGA AAG GGT TTG GAA	48
Asn Tyr Ala Met Asn Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu	
1 5 10 15	
TGG GTT GCT	57
Trp Val Ala	

(2) INFORMATION FOR SEQ ID NO:89:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 19 amino acids
- (B) TYPE: amino acid
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:89:

Asn Tyr Ala Met Asn Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu
1 5 10 15

Trp Val Ala

(2) INFORMATION FOR SEQ ID NO:90:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 57 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(ix) FEATURE:

- (A) NAME/KEY: CDS
- (B) LOCATION: 1..57

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:90:

CGC ATA AGA AGT AAA AGT AAT AAT TAT GCA ACA TTT TAT GCC GAT TCA	48
Arg Ile Arg Ser Lys Ser Asn Asn Tyr Ala Thr Phe Tyr Ala Asp Ser	
1 5 10 15	
GTG AAA GAC	57
Val Lys Asp	

(2) INFORMATION FOR SEQ ID NO:91:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 19 amino acids
 - (B) TYPE: amino acid
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:91:

Arg Ile Arg Ser Lys Ser Asn Asn Tyr Ala Thr Phe Tyr Ala Asp Ser	
1 5 10 15	
Val Lys Asp	

(2) INFORMATION FOR SEQ ID NO:92:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 96 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(ix) FEATURE:

- (A) NAME/KEY: CDS
- (B) LOCATION: 1..96

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:92:

AGG	TTC	ACC	ATC	TCC	AGA	GAT	GAT	TCA	CAA	AGC	ATG	CTC	TAT	CTG	CAA	48
Arg	Phe	Thr	Ile	Ser	Arg	Asp	Asp	Ser	Gln	Ser	Met	Leu	Tyr	Leu	Gln	
1				5				10						15		
ATG	AAC	AAC	TTG	AAA	ACT	GAG	GAC	ACA	GCC	ATG	TAT	TAC	TGT	GTG	AGA	96
Met	Asn	Asn	Leu	Lys	Thr	Glu	Asp	Thr	Ala	Met	Tyr	Tyr	Cys	Val	Arg	
			20					25					30			

(2) INFORMATION FOR SEQ ID NO:93:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 32 amino acids
- (B) TYPE: amino acid
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:93:

Arg	Phe	Thr	Ile	Ser	Arg	Asp	Asp	Ser	Gln	Ser	Met	Leu	Tyr	Leu	Gln
1				5				10						15	
Met	Asn	Asn	Leu	Lys	Thr	Glu	Asp	Thr	Ala	Met	Tyr	Tyr	Cys	Val	Arg
			20					25					30		

(2) INFORMATION FOR SEQ ID NO:94:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 69 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

- (ix) FEATURE:
 (A) NAME/KEY: CDS
 (B) LOCATION: 1..69

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:94:

CGG GGG GCT TCA GGG ATT GAC TAT GCT ATG GAC TAC TGG GGT CAA GGA 48
Arg Gly Ala Ser Gly Ile Asp Tyr Ala Met Asp Tyr Trp Gly Gln Gly
 1 5 10 15

ACC TCA CTC ACC GTC TCC TCA 69
Thr Ser Leu Thr Val Ser Ser
 20

(2) INFORMATION FOR SEQ ID NO:95:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 23 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:95:

Arg Gly Ala Ser Gly Ile Asp Tyr Ala Met Asp Tyr Trp Gly Gln Gly
 1 5 10 15

Thr Ser Leu Thr Val Ser Ser
 20

	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	2022	2023	2024	2025	2026	2027	2028	2029	2030	2031	2032	2033	2034	2035	2036	2037	2038	2039	2040	2041	2042	2043	2044	2045	2046	2047	2048	2049	2050	2051	2052	2053	2054	2055	2056	2057	2058	2059	2060	2061	2062	2063	2064	2065	2066	2067	2068	2069	2070	2071	2072	2073	2074	2075	2076	2077	2078	2079	2080	2081	2082	2083	2084	2085	2086	2087	2088	2089	2090	2091	2092	2093	2094	2095	2096	2097	2098	2099	2100	2101	2102	2103	2104	2105	2106	2107	2108	2109	2110	2111	2112	2113	2114	2115	2116	2117	2118	2119	2120	2121	2122	2123	2124	2125	2126	2127	2128	2129	2130	2131	2132	2133	2134	2135	2136	2137	2138	2139	2140	2141	2142	2143	2144	2145	2146	2147	2148	2149	2150	2151	2152	2153	2154	2155	2156	2157	2158	2159	2160	2161	2162	2163	2164	2165	2166	2167	2168	2169	2170	2171	2172	2173	2174	2175	2176	2177	2178	2179	2180	2181	2182	2183	2184	2185	2186	2187	2188	2189	2190	2191	2192	2193	2194	2195	2196	2197	2198	2199	2200	2201	2202	2203	2204	2205	2206	2207	2208	2209	2210	2211	2212	2213	2214	2215	2216	2217	2218	2219	2220	2221	2222	2223	2224	2225	2226	2227	2228	2229	2230	2231	2232	2233	2234	2235	2236	2237	2238	2239	2240	2241	2242	2243	2244	2245	2246	2247	2248	2249	2250	2251	2252	2253	2254	2255	2256	2257	2258	2259	2260	2261	2262	2263	2264	2265	2266	2267	2268	2269	2270	2271	2272	2273	2274	2275	2276	2277	2278	2279	2280	2281	2282	2283	2284	2285	2286	2287	2288	2289	2290	2291	2292	2293	2294	2295	2296	2297	2298	2299	2300	2301	2302	2303	2304	2305	2306	2307	2308	2309	2310	2311	2312	2313	2314	2315	2316	2317	2318	2319	2320	2321	2322	2323	2324	2325	2326	2327	2328	2329	2330	2331	2332	2333	2334	2335	2336	2337	2338	2339	2340	2341	2342	2343	2344	2345	2346	2347	2348	2349	2350	2351	2352	2353	2354	2355	2356	2357	2358	2359	2360	2361	2362	2363	2364	2365	2366	2367	2368	2369	2370	2371	2372	2373	2374	2375	2376	2377	2378	2379	2380	2381	2382	2383	2384	2385	2386	2387	2388	2389	2390	2391	2392	2393	2394	2395	2396	2397	2398	2399	2400	2401	2402	2403	2404	2405	2406	2407	2408	2409	2410	2411	2412	2413	2414	2415	2416	2417	2418	2419	2420	2421	2422	2423	2424	2425	2426	2427	2428	2429	2430	2431	2432	2
--	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	---

(A) LENGTH: 69 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ix) FEATURE:

(A) NAME/KEY: CDS
(B) LOCATION: 1..69

[illegible]

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 23 amino acids
- (B) TYPE: amino acid
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:97:

Gln Ile Val Leu Ser Gln Ser Pro Ala Ile Leu Ser Ala Ser Pro Gly
1 5 10 15
Glu Lys Val Thr Met Thr Cys
20

(2) INFORMATION FOR SEQ ID NO:98:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 30 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

- (ix) FEATURE:
 (A) NAME/KEY: CDS
 (B) LOCATION: 1..30

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:98:

AGG GCC AGC TCA AGT GTA AAT TAC ATG CAC
Arg Ala Ser Ser Ser Val Asn Tyr Met His
 1 5 10

30

(2) INFORMATION FOR SEQ ID NO:99:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 10 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:99:

Arg Ala Ser Ser Ser Val Asn Tyr Met His
 1 5 10

(2) INFORMATION FOR SEQ ID NO:100:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 66 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

- (ix) FEATURE:
 (A) NAME/KEY: CDS
 (B) LOCATION: 1..66

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:100:

TGG TAC CAG CAG AAG CCA GGA TCC TCC CCC AAA CCC TGG ATT TCT GCC 48
Trp Tyr Gln Gln Lys Pro Gly Ser Ser Pro Lys Pro Trp Ile Ser Ala
1 5 10 15
ACA TCC AAC CTG GCT TCT 66
Thr Ser Asn Leu Ala Ser
20

(2) INFORMATION FOR SEQ ID NO:101:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 22 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:101:

Trp Tyr Gln Gln Lys Pro Gly Ser Ser Pro Lys Pro Trp Ile Ser Ala
1 5 10 15
Thr Ser Asn Leu Ala Ser
20

[illegible]

(A) LENGTH: 96 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(ix) FEATURE:

(A) NAME/KEY: CDS
(B) LOCATION: 1..96

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:102:

GGA	GTC	CCT	GCT	CGC	TTC	AGT	GGC	AGT	GGG	TCT	GGG	ACC	TCT	TAC	TCT	48
Gly	Val	Pro	Ala	Arg	Phe	Ser	Gly	Ser	Gly	Ser	Gly	Thr	Ser	Tyr	Ser	
1				5					10					15		
CTC	ACA	ATC	AGC	AGA	GTG	GAG	GCT	GAA	GAT	GCT	GCC	ACT	TAT	TAC	TGC	96
Leu	Thr	Ile	Ser	Arg	Val	Glu	Ala	Glu	Asp	Ala	Ala	Thr	Tyr	Tyr	Cys	
			20					25					30			

(2) INFORMATION FOR SEQ ID NO:103:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 32 amino acids
(B) TYPE: amino acid
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:103:

Gly Val Pro Ala Arg Phe Ser Gly Ser Gly Ser Gly Thr Ser Tyr Ser
1 5 10 15
Leu Thr Ile Ser Arg Val Glu Ala Glu Asp Ala Ala Thr Tyr Tyr Cys
20 25 30

(2) INFORMATION FOR SEQ ID NO:104:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 57 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

- (ix) FEATURE:
 (A) NAME/KEY: CDS
 (B) LOCATION: 1..57

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:104:

CAG CAG TGG AGT AGT AAC CCA CCC ACG TTC GGA GGG GGG ACC ATG CTG	48
Gln Gln Trp Ser Ser Asn Pro Pro Thr Phe Gly Gly Gly Thr Met Leu	
1 5 10 15	
GAA ATA AGA	57
Glu Ile Arg	

(2) INFORMATION FOR SEQ ID NO:105:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 19 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:105:

Gln Gln Trp Ser Ser Asn Pro Pro Thr Phe Gly Gly Gly Thr Met Leu
1 5 10 15
Glu Ile Arg